

## Glassy-winged sharpshooter

### What is the glassy-winged sharpshooter?

The glassy-winged sharpshooter (GWSS) (*Homalodisca vitripennis*) is a xylem feeding leafhopper that causes direct damage to citrus plants through its feeding activities. However, the greatest threat is its potential to vector the bacterium *Xylella fastidiosa* subsp. *pauca* which causes citrus variegated chlorosis. Over 100 plant species are hosts of GWSS including many commercial hosts like grape, almond and peach as well as several ornamentals. These hosts would be severely threatened if GWSS became established in Australia, particularly if it is carrying *X. fastidiosa*.



Adult GWSS showing white spots of excrement



Adult GWSS on leaf surface



Adult GWSS on stem

### What does it look like?

Adult GWSS are easily visible with the naked eye, measuring about 12-14 mm long with a dark brown to black colouring and a lighter underside. The upper parts of the head and back are dotted with ivory or yellowish spots and their wings are partly transparent with reddish veins. Watery excrement often collects on the sides of the insect, appearing as large white spots.

Clutches of up to 27 eggs are laid on the underside of leaves of citrus plants in a side-by-side arrangement and are dusted with a layer of whitish powder. Following hatching they change in appearance from green water blisters to tan or brown scars on the leaves. The immature nymphs that hatch from the eggs are wingless.

### What can it be confused with?

Australian leafhoppers from the Auchenorrhyncha suborder share some features with GWSS including larger size, brown colouration and large head with prominent eyes.



## What should I look for?

GWSS excretes copious amounts of liquid that can make leaves, stems and fruit appear white washed when dry. Look for egg masses that are usually laid into recently expanded foliage. Older foliage will contain the distinctive tan or brown scars left after the eggs have hatched. If *X. fastidiosa* enters Australia with the GWSS, the symptoms of citrus variegated chlorosis, such as leaf yellowing, brown lesions on the underside of leaves, leaf drop and small, hard fruit may be observed.



GWSS egg mass on leaf ready to hatch

## How does it spread?

Adult GWSS are strong fliers and can move rapidly from plant to plant. Nymphs are wingless but can spread throughout the orchard by walking and jumping through the canopy or along the ground to a new host. Most rapid and long distance movement occurs through the transportation of viable egg masses on nursery stock of either crop or ornamental plants.

## Where is it now?

GWSS is found in eastern and western USA, Mexico, Tahiti, French Polynesia and Hawaii. It has most recently spread to Easter Island and the Cook Islands.



Yellowing of leaves in citrus infected with citrus variegated chlorosis

## How can I protect my orchard from the Glassy-winged sharpshooter?

Check your orchard frequently for the presence of new pests and unusual symptoms. Make sure you are familiar with common citrus pests so you can tell if you see something different. Keep records of anything unusual and ensure that all staff and visitors adhere to orchard biosecurity and hygiene practices.

**If you see anything unusual, call the Exotic Plant Pest Hotline**



**Disclaimer:** The material in this publication is for general information only and no person should act, or fail to act on the basis of this material without first obtaining professional advice. Plant Health Australia and all persons acting for Plant Health Australia expressly disclaim liability with respect to anything done in reliance on this publication.

For more information visit [www.planthealthaustralia.com.au](http://www.planthealthaustralia.com.au)