

Liriomyza huidobrensis Contingency Plan

Prepared for Horticulture Innovation Australia, as part of Project MT16004 (*RD&E program for control, eradication and preparedness for vegetable leafminer*)

Rohan Burgess¹, Dr. Peter Ridland², Dr. Elia Pirtle³

¹ Plant Health Australia

² The University of Melbourne

³ Cesar Australia

December 7, 2020

This resource has been funded by Horticulture Innovation, through the Project MT16004 (*RD&E program for control, eradication and preparedness for vegetable leafminer*), using multiple research and development levies and contributions from the Australian Government. Hort Innovation is the grower-owned, not-for-profit research and development corporation for Australian horticulture.

**Hort
Innovation**



CONTINGENCY PLAN

SERPENTINE LEAFMINER

(*LIRIOMYZA HUIDOBRENSIS*)



Merle Shepard, Gerald R. Carner, and P.A.C. Ooi, *Insects and their Natural Enemies*
Associated with Vegetables and Soybean in Southeast Asia, Bugwood.org



Central Science Laboratory, Harpenden, British Crown,
Bugwood.org

December 2020

This resource has been funded by Horticulture Innovation, through the Project MT16004 (RD&E program for control, eradication and preparedness for vegetable leafminer), using multiple research and development levies and contributions from the Australian Government. Hort Innovation is the grower-owned, not-for-profit research and development corporation for Australian horticulture.

**Hort
Innovation**



This Contingency Plan has been authored by Rohan Burgess (Plant Health Australia), Dr. Peter Ridland (The University of Melbourne) and Dr. Elia Pirtle (Cesar Australia), with contributions from Dr. Sharyn Taylor (Plant Health Australia), Dr. James Maino (Cesar Australia), and Dr. Paul Umina (Cesar Australia).

Disclaimer

The scientific and technical content of this document is current to the date published and all efforts have been made to obtain relevant and published information on these pests. New information will be included as it becomes available, or when the document is reviewed. The material contained in this publication is produced for general information only. It is not intended as professional advice on any particular matter. No person should act or fail to act on the basis of any material contained in this publication without first obtaining specific, independent professional advice. Plant Health Australia and all persons acting for Plant Health Australia in preparing this publication, expressly disclaim all and any liability to any persons in respect of anything done by any such person in reliance, whether in whole or in part, on this publication. The views expressed in this publication are not necessarily those of Plant Health Australia.

Further information

For further information regarding this contingency plan, contact Plant Health Australia through the details below.



Address:	Level 1, 1 Phipps Close DEAKIN ACT 2600
Phone:	+61 2 6215 7700
Fax:	+61 2 6260 4321
Email:	mailto:biosecurity@phau.com.au
Website:	www.planthealthaustralia.com.au

An electronic copy of this plan is available from the web site listed above.

© Plant Health Australia Limited 2020

Copyright in this publication is owned by Plant Health Australia Limited, except when content has been provided by other contributors, in which case copyright may be owned by another person. With the exception of any material protected by a trade mark, this publication is licensed under a [Creative Commons Attribution-No Derivs 3.0 Australia licence](https://creativecommons.org/licenses/by-nd/3.0/). Any use of this publication, other than as authorised under this licence or copyright law, is prohibited.



<http://creativecommons.org/licenses/by-nd/3.0/> - This details the relevant licence conditions, including the full legal code. This licence allows for redistribution, commercial and non-commercial, as long as it is passed along unchanged and in whole, with credit to Plant Health Australia (as below).

In referencing this document, the preferred citation is:

Plant Health Australia Limited (Version 1, 2020) Contingency Plan for Serpentine Leafminer (*Liriomyza huidobrensis*).

1. Purpose and background of this contingency plan	6
2. Pest details.....	7
2.1 Background and impact of pest	7
2.2 Life cycle.....	7
2.3 Host range.....	8
2.4 Signs and symptoms.....	8
2.5 Dispersal.....	10
2.6 Current geographic distribution.....	10
2.7 Risk of establishment within Australia	13
2.8 Risk pathways within Australia	14
2.8.1 Risk pathways – international and domestic.....	14
2.8.2 Predicting spread potential.....	19
2.9 Risk of economic impact within Australia.....	20
3. Diagnostic information	23
3.1 Diagnostic considerations.....	23
4. Surveillance and collection of samples.....	25
4.1 Surveillance.....	25
4.1.1 Technical information for planning surveys	25
4.1.2 Delimiting surveys in the event of an incursion into new areas	28
4.1.3 Activities for general surveillance immediately following a detection.....	30
4.2 Collection and treatment of samples	30
4.2.1 If collecting for morphological diagnosis.....	30
4.2.2 If collecting for molecular diagnosis	31
5. Immediate response to detection	34
5.1 Tracing	34
5.2 Quarantine and movement controls.....	34
5.3 Destruction strategy.....	34
5.3.1 Priorities and considerations	34
5.3.2 Decontamination of tools and equipment.....	35
5.3.3 Chemical control	35
5.3.4 Physical control options	40
5.3.5 Biological control	40
5.4 Decision support for eradication.....	41
6. Emergency Plant Pest Response Deed and PLANTPLAN.....	47
6.1 Emergency Plant Pest Response Deed.....	47
6.2 PLANTPLAN	48
7. References.....	49
8. Appendices.....	55
Appendix 1: Known hosts of serpentine leafminer	55
Number of host records for <i>L. huidobrensis</i> within plant families.....	55
Commercial, ornamental and non-cultivated plants for which comprehensive evidence of lifecycle completion in the field for the serpentine leafminer (<i>Liriomyza huidobrensis</i>) has been provided within the scientific literature.....	57

Commercial, ornamental, and non-cultivated plants for which only partial evidence of lifecycle completion in the field for the serpentine leafminer (<i>Liriomyza huidobrensis</i>) has been provided within the scientific literature.....	72
Commercial, ornamental and non-cultivated plants with unverified records of lifecycle completion in the field for the serpentine leafminer (<i>Liriomyza huidobrensis</i>).....	89
References	92
Appendix 2: Leaf mining damage caused by established species	97
References	109
Appendix 3: Permits for the control of serpentine leafminer and other <i>Liriomyza</i> spp. on different host crops.....	110
Appendix 4: Additional impact prediction outputs.....	127

1. PURPOSE AND BACKGROUND OF THIS CONTINGENCY PLAN

The serpentine leafminer (*Liriomyza huidobrensis*) is an exotic leafminer that affects a wide range of horticultural industries, including the vegetable and melon industries. It is currently widespread overseas and is present in nearby countries such as Indonesia.

This Contingency Plan provides background information on serpentine leafminer to assist in determining the requirements for the initial response to a detection of this species in Australia. Only key information to support an immediate response is provided in this document.

Additional information can be found in the following supporting material:

- Awareness material such as the fact sheets from Plant Health Australia (PHA), AUSVEG, and commonwealth, state and territory jurisdictions.
 - www.planthealthaustralia.com.au/pests/pea-leaf-miner/
 - www.agriculture.gov.au/pests-diseases-weeds/plant/leaf-miner
 - www.dpi.nsw.gov.au/biosecurity/plant/insect-pests-and-plant-diseases/exotic-leaf-miners
 - <https://ausveg.com.au/biosecurity-agrichemical/biosecurity/mt16004/>
- Overseas websites with additional information
 - <https://gd.eppo.int/taxon/LIRIHU>
 - www.cabi.org/isc/datasheet/30956
 - <https://cipotato.org/riskatlasforafrica/liriomyza-huidobrensis/>
 - <https://ufdc.ufl.edu/IR00004169/00001>
 - <http://ipm.ucanr.edu/PMG/r280300911.html>

2. PEST DETAILS

Preferred common name:	Serpentine leafminer
Other common names	Pea Leafminer; South American leafminer
Scientific name:	<i>Liriomyza huidobrensis</i> (Blanchard, 1926)
Synonyms:	<i>Agromyza huidobrensis</i> <i>Liriomyza cucumifoliae</i> <i>Liriomyza decora</i> <i>Liriomyza dianthi</i>
Taxonomic position:	Class: Insecta Order: Diptera Family: Agromyzidae Genus: <i>Liriomyza</i> Species: <i>Liriomyza huidobrensis</i>

2.1 Background and impact of pest

The Agromyzidae are a well-known group of small, morphologically similar flies whose larvae feed internally on plants, often as leaf and stem miners. Nearly all species are very host-specific, but a few highly polyphagous species have become important pests of agriculture and horticulture in many parts of the world.

The serpentine leafminer is one such species and is of concern as it has a wide host range including many horticulture crops in which it can cause significant yield losses and quality reductions (see Appendix 1). The serpentine leafminer is widespread overseas being present in North and South America, Asia, Africa, and Europe.

2.2 Life cycle

The lifecycle of the serpentine leafminer consists of an egg inserted by a female just under the surface of the leaf of a host plant. The egg hatches in 2-5 days. The larvae then begin to feed within the leaf creating tunnels or mines that get larger as the larvae moult and mature. After passing through three larval stages in 4-7 days the larva leaves the plant to form a puparium in the soil underneath the host plant. After 7-14 days (at 20°C and 30°C) an adult emerges and begins to reproduce (CABI and EPPO date unknown). The time taken to complete each life stage vary depending on host and temperature. With cooler temperatures resulting in longer development times. For example, Lanzoni et al., (2002) studied the effect of temperature on serpentine leafminer on common bean at 15, 20, 25 and 30°C, their results found total development times ranged from 43.6 days at 15°C to 16.1 days at 25°C, no adults emerged at 30°C. Similarly, Mujica et al., (2017) investigated serpentine leafminer development on faba bean and found development took between 65.62 days at 10°C to 13.48 days at 30°C, and that the lifecycle was not completed at 32 or 35°C.

The serpentine leafminer is reported to be able to survive winter temperatures as low as -11.5°C as pupae and can complete its lifecycle at temperatures between 5.7 and 30°C (EFSA Panel on Plant Health 2012).

2.3 Host range

Serpentine leafminer is a polyphagous pest of many agricultural and ornamental hosts (see Appendix 1) It has been recorded from over 250 plant species representing 50 plant families.

Important agricultural hosts include pulses, oilseeds, vegetables, herbs, cereals, and a wide range of ornamental species such as chrysanthemum as well as weed and pasture species.

A detailed host list is included in Appendix 1.

2.4 Signs and symptoms

Adult flies are small (1.3-2.3 mm long with a wing length of 1.5-2.2 mm (Okoth et al., 2014)) with a shiny black mesonotum and yellow markings on the head and (Figure 1).

Female flies use their ovipositor to puncture the leaves of the host plants causing wounds which serve as sites for feeding (by both males and females) or oviposition. Feeding punctures appear as white speckles between 0.13 and 0.15 mm in diameter. Oviposition punctures are usually smaller (0.05 mm) and are more uniformly round. However, the appearance of the punctures does not differ between *Liriomyza* species so cannot be used to separate species.

Leaf mines are pale coloured with black and brown areas. Mines are created by the feeding larvae and are the most obvious symptom of infestations of *Liriomyza* spp. (Figure 2). Mines can be linear or serpentine and increase in width along their length as the larvae matures.

Factsheets with more information on what to look for in the field can be found at the following websites:

- www.planthealthaustralia.com.au/pests/pea-leaf-miner/
- <https://ufdc.ufl.edu/IR00004169/00001>
- <http://ipm.ucanr.edu/PMG/r280300911.html>
- <https://ausveg.com.au/biosecurity-agrichemical/biosecurity/mt16004/>



Figure 1 Adult serpentine leafminer. Source: Central Science Laboratory, Harpenden, British Crown, Bugwood.org



Figure 2 Leaf mines on chrysanthemum. Source: National Plant Protection Organization, the Netherlands, Bugwood.org

2.5 Dispersal

Typically, *Liriomyza* leafminers are considered to have invaded countries via movement of infested plants (generally on ornamentals, but also some horticultural crops such as pea pods). While fully formed mines should be readily visible, signs of early infestations are much less obvious and are easily overlooked.

Agromyzid flies are considered as “moderate fliers” (Yoshimoto and Gressitt 1964) and in agricultural situations, the flies tend to remain close to their target crops, often only moving very short distances between host plants (Fenoglio et al., 2019). They also have the potential to be spread over large distances through wind dispersal. Flight mill experiments on the related *L. sativae* have shown that leafminers can fly up to 8.22 km but that the average flight distances was <1.0 km at 18-36°C (Lei et al., 2002). Similar flight ability is expected for serpentine leafminer.

Serpentine leafminer adults have been shown to disperse on average 46 m within 24 hours via a release recapture experiment (Fenoglio et al 2019). Similarly, *Liriomyza trifolii* individuals within glasshouses have been shown to disperse on average 26 m but up to 100 m from original infestation points within 24 hours (Jones and Parella 1986) and in a field study were generally not found more than 50 m away from infested glasshouses (Ozawa et al 1999). *Agromyza frontella*, a related agromyzid leafminer pest of lucerne with similar flight dynamics, was observed to spread at a rate of about 100 km per year (or averaging about 0.3 kms per day) during an incursion into the Midwestern United States (Venette et al., 1980). There is also some evidence of long distance dispersal of Agromyzids via wind, with Agromyzid species reportedly trapped as high as 900 meters in the air column (Hardy and Milne 1938; Glick 1960; White 1970), and over oceans (Cheng 1976, Yoshimoto and Gressitt 1964).

However, windborne dispersal is likely to be a minor pathway of leafminer dispersal in Australia, and human-mediated pathways should be prioritised, particularly given the accumulation of research pointing to this as a major contributor to the movement of invasive species (Capinha et al. 2015; Wichmann et al. 2009). Such pathways include transport of contaminated produce, or “hitch-hiking” adults associated with the large volume of human movement.

Since the 1980s the serpentine leafminer has spread from South America into Central America, North America, Europe, Africa and Asia through the movement of plant material (cuttings, cut flowers) (CABI and EPPO date unknown). This suggests that there is the potential for it to spread rapidly to new regions. Plant material should be considered as a potential pathway for the introduction/spread of this pest. The introduction of plant material from overseas is regulated, however illegal movement of plant material poses a potential risk for the entry of the pest into the country.

2.6 Current geographic distribution

The serpentine leafminer is thought to have originated in South America and since spread to North and Central America, Asia, Africa and Europe.

The current distribution of this pest is presented in Table 1. Note, serpentine leafminer has also been found at various times in a number of European countries including Bulgaria, Denmark, Finland, Hungary, Ireland, Lithuania, Slovenia, United Kingdom, but has since been eradicated and are no longer present in those countries (EFSA Panel on Plant Health 2012).

Table 1. Countries where serpentine leafminer is known to occur

COUNTRY/REGION	REFERENCE
Argentina	CABI and EPPO (date unknown); Maharjan et al., (2014); Milla and Reitz (2005); Scheffer and Lewis (2001); Spencer (1989); Weintraub et al., (2017)
Austria	CABI and EPPO (date unknown); EFSA Panel on Plant Health (2012); Maharjan et al., (2014)
Belgium	CABI and EPPO (date unknown); EFSA Panel on Plant Health (2012); Maharjan et al., (2014); Milla and Reitz (2005); Weintraub et al., (2017)
Belize	CABI and EPPO (date unknown); Maharjan et al., (2014); Milla and Reitz (2005)
Brazil	CABI and EPPO (date unknown); Maharjan et al., (2014); Milla and Reitz (2005); Spencer (1973); Weintraub et al., (2017)
Canada	Marin et al., (2005); Milla and Reitz (2005); Scheffer et al., (2001); Weintraub et al., (2017)
Chile	CABI and EPPO (date unknown); Maharjan et al., (2014); Milla and Reitz (2005); Weintraub et al., (2017); Spencer (1973)
China	He et al., (2002); Maharjan et al., (2014); Milla and Reitz (2005); Weintraub et al., (2017)
Colombia	CABI and EPPO (date unknown); Maharjan et al., (2014); Milla and Reitz (2005); Spencer (1973); Weintraub et al., (2017)
Comoros	Maharjan et al., (2014)
Costa Rica	CABI and EPPO (date unknown); Maharjan et al., (2014); Milla and Reitz (2005); Scheffer and Lewis (2001); Spencer (1983); Weintraub et al., (2017)
Croatia	Maharjan et al., (2014)
Cyprus	CABI and EPPO (date unknown); EFSA Panel on Plant Health (2012); Maharjan et al., (2014)
Czech Republic	CABI and EPPO (date unknown); EFSA Panel on Plant Health (2012); Maharjan et al., (2014)
Dominican Republic	CABI and EPPO (date unknown); Maharjan et al., (2014); Milla and Reitz (2005); Spencer (1989)
Ecuador	Maharjan et al., (2014); Scheffer et al., (2001); Weintraub et al., (2017)
El Salvador	CABI and EPPO (date unknown); Maharjan et al., (2014); Milla and Reitz (2005)
France	CABI and EPPO (date unknown); EFSA Panel on Plant Health (2012); Maharjan et al., (2014); Trouvé et al., (1991); Weintraub et al., (2017)
French Guiana	Maharjan et al., (2014)
Germany	EFSA Panel on Plant Health (2012); Maharjan et al., (2014); Weintraub et al., (2017)
Greece	EFSA Panel on Plant Health (2012); Maharjan et al., (2014); Weintraub et al., (2017)
Guadeloupe	CABI and EPPO (date unknown); Maharjan et al., (2014); Milla and Reitz

COUNTRY/REGION	REFERENCE
	(2005); Spencer et al., (1992)
Guatemala	CABI and EPPO (date unknown); Maharjan et al., (2014); Milla and Reitz (2005); Scheffer et al., (2001); Weintraub et al., (2017)
Honduras	CABI and EPPO (date unknown); Maharjan et al., (2014); Milla and Reitz (2005)
India	CABI and EPPO (date unknown); Maharjan et al., (2014); Singh and Singh (2013)
Indonesia	Maharjan et al., (2014); Milla and Reitz (2005); Scheffer et al., (2001); Shepard and Braun (1998); Weintraub et al., (2017)
Israel	CABI and EPPO (date unknown); Maharjan et al., (2014); Milla and Reitz (2005); Scheffer et al., (2001); Weintraub et al., (2017); Weintraub and Horowitz (1995)
Italy	CABI and EPPO (date unknown); Suss (1991); EFSA Panel on Plant Health (2012); Maharjan et al., (2014); Weintraub et al., (2017)
Japan	Weintraub et al., (2017)
Jordon	Maharjan et al., (2014)
Kenya	Guantai et al., (2015); Maharjan et al., (2014); Milla and Reitz (2005); Weintraub et al., (2017);
Lebanon	Maharjan et al., (2014); Weintraub et al., (2017)
Malaysia	Maharjan et al., (2014); Weintraub et al., (2017)
Malta	CABI and EPPO (date unknown); EFSA Panel on Plant Health (2012); Maharjan et al., (2014)
Mauritius	CABI and EPPO (date unknown); Maharjan et al., (2014); Weintraub et al., (2017)
Mexico	Maharjan et al., (2014)
Morocco	Hanafi (2005); Maharjan et al., (2014); Weintraub et al., (2017)
Nepal	Maharjan et al., (2014)
Netherlands	CABI and EPPO (date unknown); EFSA Panel on Plant Health (2012); Maharjan et al., (2014); Milla and Reitz (2005); Weintraub et al., (2017)
Nicaragua	CABI and EPPO (date unknown); Maharjan et al., (2014); Milla and Reitz (2005)
North Korea	Maharjan et al., (2014)
Norway	Maharjan et al., (2014); Weintraub et al., (2017)
Panama	CABI and EPPO (date unknown); Maharjan et al., (2014); Milla and Reitz (2005)
Peru	CABI and EPPO (date unknown); Maharjan et al., (2014); Milla and Reitz (2005); Spencer (1973); Weintraub et al., (2017)
Philippines	Weintraub et al., (2017)
Poland	EFSA Panel on Plant Health (2012); Maharjan et al., (2014)
Portugal	CABI and EPPO (date unknown); EFSA Panel on Plant Health (2012);

COUNTRY/REGION	REFERENCE
	Maharjan et al., (2014); Weintraub et al., (2017)
Réunion	CABI and EPPO (date unknown); Maharjan et al., (2014); Vercambre and De Crozals (1993); Weintraub et al., (2017)
Saudi Arabia	Dawah and Deeming (2002)
Serbia	Dobrosavljevic et al., (2017)
Seychelles	Maharjan et al., (2014)
Singapore	Maharjan et al., (2014)
South Africa	Maharjan et al., (2014); Milla and Reitz (2005); Scheffer et al., (2001); Weintraub et al., (2017)
South Korea	Maharjan et al., (2014); Weintraub et al., (2017)
Spain	CABI and EPPO (date unknown); EFSA Panel on Plant Health (2012); Maharjan et al., (2014); Weintraub et al., (2017)
Sri Lanka	Maharjan et al., (2014); Milla and Reitz (2005); Scheffer et al., (2001); Weintraub et al., (2017)
Sweden	Maharjan et al., (2014)
Switzerland	Maharjan et al., (2014)
Syria	Maharjan et al., (2014)
Taiwan	Milla and Reitz (2005); Shiao and Wu (2000); Weintraub et al., (2017)
Thailand	CABI and EPPO (date unknown); Maharjan et al., (2014); Weintraub et al., (2017)
Turkey	Maharjan et al., (2014); Weintraub et al., (2017)
United States of America (including Hawaii)	CABI and EPPO (date unknown); Maharjan et al., (2014); Scheffer and Lewis (2001); Spencer (1973); Spencer (1989); Weintraub et al., (2017)
Uruguay	Maharjan et al., (2014)
Venezuela	CABI and EPPO (date unknown); Maharjan et al., (2014); Milla and Reitz (2005); Spencer (1973);
Vietnam	Maharjan et al., (2014); Weintraub et al., (2017)
Zimbabwe	Weintraub et al., (2017)

2.7 Risk of establishment within Australia

CLIMEX models are provided in Jovicich (2009). These found that the serpentine leafminer is most likely to establish along the eastern, southern and southwestern coastline of Australia and Tasmania.

Refer to: <https://ausveg.com.au/app/data/technical-insights/docs/VG06113.pdf> for further details.

Similarly, Weintraub et al., (2017) suggest serpentine leafminer may likely establish in parts of Australia that have mild winters (i.e. do not receive snow) and have summer highs of 30-35°C.

Cesar Australia has created a model of the potential geographic distribution of the serpentine leafminer in Australia, the results of which are summarised below (Figure 3). A copy of the relevant




Figure 3 Aggregated yearly establishment potential of *L. huidobrensis* as the number of months across the year with increasing population sizes) based on temperature and moisture constraints (left) and predicted dominant stressors based on the highest mortality rate from desiccation, cold, and heat stress across the year (right).

2.8 Risk pathways within Australia

2.8.1 Risk pathways – international and domestic

This species could potentially enter new areas by hitchhiking on goods, aircraft, vehicles, or via the movement of plant material (DAWR 2016). Table 2 lists some of the potential international risk pathways and control measures in place. These are also be broadly applicable to other *Liriomyza* species.

Table 2. International risk pathways for the spread of the pest and control measures

RISK PATHWAY	DESCRIPTION	CONTROL MEASURES
Cut flowers on regulated imports	Infested cut flowers moved to residential dwellings that contain suitable hosts and the insect transfers to a host.	Border inspections
Infested plant material on passenger or mail	Infested plant material moved to residential dwellings that contain	Border inspections

RISK PATHWAY	DESCRIPTION	CONTROL MEASURES
pathways	suitable hosts and the insect transfers to a host.	
Natural dispersal from Papua New Guinea (PNG)	Ongoing spread from PNG to Torres Strait and/or Far North Queensland via traditional movements, winds or flight.	Ongoing surveillance via the Northern Australia Quarantine Strategy Domestic quarantine
Infested plant material	Plant material or pupae in soil spread to new areas in baggage, vehicles or camping equipment.	Domestic quarantine

Additionally, as the larvae of *Liriomyza* spp. leave the plant to pupate, pupae may also be spread with crop debris or soil associated with infested areas. Table 3 lists some of the potential risk pathways and potential domestic control measures to reduce the spread of serpentine leafminer. These will also be broadly applicable to other *Liriomyza* species.

Table 3. Risk pathways for the spread of the pest and potential domestic control measures

RISK PATHWAY	DESCRIPTION	POTENTIAL DOMESTIC CONTROL MEASURES
Cut flowers (host plants)	<p>Cut flowers that are a host of serpentine leafminer (see Appendix 1) pose a significant risk. The life of cut flowers is long enough to allow the completion of the pest's lifecycle (CABI and EPPO, date unknown). Eggs may be present within leaves but are never visible.</p> <p>Larvae may be present and visible within leaves.</p> <p>Hitchhiking adults and pupae may be transported on cut flowers.</p>	<p>The application of systemic insecticides may be required before allowing cut flowers grown in an infested area to be moved outside the infested area. Further information on chemical control is provided in Section 5.3. 3.</p> <p>All cut flowers leaving infested areas will also need to be visually inspected for signs of leafminers before leaving the area.</p>
Cut flowers (non-host plants)	<p>Cut flowers that are not a host of serpentine leafminer poses significantly less risk for the spread of this pest, but is still a potential pathway, particularly if the non-host cut flowers are transported with or spent time in close proximity with host cut flowers.</p> <p>Eggs, larvae and pupae will not spread with non-host stock.</p> <p>Adults may potentially spread with non-host stock as a hitchhiker.</p>	<p>To reduce the risk of spread, cut flowers that are not a host plant for vegetable leafminer should be inspected and/or treated as a precaution if grown in an area known to be infested with serpentine leafminer.</p>
Nursery Stock (containerized host plants)	<p>Nursery stock that is a host plant for serpentine leafminer (see Appendix 1) poses a significant risk for the spread of this pest.</p> <p>Containerized nursery stock can carry all life stages of the pest.</p> <p>Eggs may be present within leaves but are never visible.</p> <p>Larvae may be present and visible within leaves.</p> <p>Pupae may also be present in the soil of containerized stock. Larvae emerge from the leaf and pupate in the soil under host plants meaning potting mix/soil can also potentially spread the pest. Larvae very occasionally attach to leaves or surrounding non-soil surfaces to pupate.</p> <p>Adults may potentially spread with containerized nursery stock as a hitchhiker.</p>	<p>Visual inspection of containerised nursery stock for the presence of adults resting on leaves, larvae creating leaf mines and pupae in the soil (see Figure 2, however pupae will be very hard to detect as they are mixed into the substrate) or attached to leaves and container surfaces.</p> <p>To reduce the risk of spread, containerised nursery stock that is a host plant for vegetable leafminer should not be moved from infested to non-infested areas without being treated with an appropriate systemic insecticide. Treatment of growing media using appropriate pesticides (eg Cyromazine-PER83506) should also be considered. Further information on chemical control is provided in Section 5.3.3.</p>
Nursery Stock	Nursery stock that is a host plant for	Visual inspection of bare rooted

RISK PATHWAY	DESCRIPTION	POTENTIAL DOMESTIC CONTROL MEASURES
(bare rooted host plants)	<p>serpentine leafminer (see Appendix 1) poses a significant risk for the spread of this pest.</p> <p>Bare rooted nursery stock can carry all life stages of the pest.</p> <p>Eggs may be present within leaves but are never visible.</p> <p>Larvae may be present and visible within leaves.</p> <p>Pupae are less likely to be present in bare rooted stock than containerized stock. However, upon emergence, if larvae are unable to find a substrate to bury into, they will attach to any surface available (including tray walls or plant leaves) meaning pupae can potentially still spread via bare rooted stock.</p> <p>Adults may potentially spread with bare rooted nursery stock as a hitchhiker.</p>	<p>nursery stock for the presence of adults resting on leaves, larvae creating leaf mines and pupae attached to leaves or surfaces surrounding the roots (see Figure 2).</p> <p>To reduce the risk of spread, bare rooted nursery stock that is a host plant for vegetable leafminer should not be moved from infested to non-infested areas without being treated with an appropriate systemic insecticide. Further information on chemical control is provided in Section 5.3.3.</p>
Nursery Stock (non-host plants)	<p>Nursery stock that is not a known host plant of serpentine leafminer poses significantly less risk for the spread of this pest, but is still a potential pathway, particularly if the non-host stock spent time in close proximity with host stock.</p> <p>Eggs, larvae and pupae will not spread with non-host stock.</p> <p>Adults may potentially spread with non-host stock as a hitchhiker.</p>	<p>Visual inspection of nursery stock for the presence of adults resting on leaves.</p> <p>To reduce the risk of spread, nursery stock that is not a host plant for vegetable leafminer should be inspected and/or treated as a precaution if grown in an area known to be infested with serpentine leafminer. Treatment of growing media using appropriate pesticides (eg Cyromazine-PER83506) may also be considered as a precaution to limit the potential for spread.</p>
Non-	People and small businesses	Should an incursion occur effort will

RISK PATHWAY	DESCRIPTION	POTENTIAL DOMESTIC CONTROL MEASURES
commercially grown seedlings and plants	<p>selling/trading seedlings at a local level (e.g. at farmers markets, school fetes, roadside stalls, etc.) pose similar threats to those posed by nursery stock, (i.e. plants can spread eggs, larvae, pupae and potentially adults via host plants to new areas). However, being less regulated this is a difficult pathway to control and therefore potentially poses a higher risk than commercial nursery businesses.</p>	<p>be needed to identify those involved in the non-commercial trade of seedlings to reduce the risks that they pose.</p> <p>Similar to commercial nursery stock, host plants should not be moved from infested to non-infested areas without being inspected and treated with an appropriate systemic insecticide. Likewise, non-host crops grown in proximity to host plants should be inspected and/or treated as a precaution if grown in an area known to be infested with serpentine leafminer, as there is a small risk that non-host crops could potentially spread adult leafminers as hitchhikers. Treatment of growing media using appropriate pesticides (eg Cyromazine-PER83506) may also be considered.</p> <p>Further information on chemical control is provided in Section 5.3.3.</p>
Plant material including hay, plant debris and crop waste	<p>Hay, plant debris and crop waste could potentially spread pupae and adults to new areas.</p> <p>Larvae are not expected to be able to survive long after plant debris and begins to wilt and rot (once it cannot feed on the plant, the larvae must vacate the leaf to begin pupation or die), however pupae will survive plant debris and crop waste.</p> <p>Adults may also hitchhike on hay, plant debris and crop waste.</p>	<p>Visual inspection for the presence of adults, larvae and pupae should be carried out.</p> <p>To reduce the potential risk of spread, plant material should not be moved from infested to non-infested areas without being treated with an appropriate insecticide. Further information on chemical control is provided in Section 5.3.3.</p>
Soil	<p>Larvae emerge from the leaf and pupate in the soil under host plants (Capinera 2014).</p> <p>Soil therefore poses a potential risk for moving the pest to new areas.</p>	<p>Minimise potential spread by reducing the spread of soil off infested sites (e.g. by cleaning down machinery between properties).</p>
Tools, equipment, and machinery	<p>Tools, equipment, and machinery used on farm in the vicinity of host plants pose a low risk of spreading the pest, however, may transport pupae via soil deposits, or may transport adults.</p>	<p>Clean tools, equipment and machinery before moving off infested sites.</p>

RISK PATHWAY	DESCRIPTION	POTENTIAL DOMESTIC CONTROL MEASURES
Transport vehicles	Vehicles in the vicinity of host plants could allow the pest to hitchhike (in cabins etc.) to new areas (DAWR 2016).	Clean vehicles before moving off infested sites.
Conveyances (includes crates, boxes, bins, pallets)	Crates, bins, pallets, etc. that are clean of soil are unlikely to spread leafminers.	As a precaution, clean conveyances before moving between sites to reduce any risk of spread.
Seed	There are no expected pathways for leafminer spread by the movement of seed.	Not applicable.
Fruit (without leaves)	There are no expected pathways for leafminer spread by the movement of fruit (without leaves). It is expected the process of harvesting and cleaning fruit will remove leaf and soil material that could spread larvae and pupae and would cause adults to disperse rather than hitchhike with fruit to new areas.	Not applicable.
Fresh leafy or green vegetables (including peas)	Mines can occur on leaves of a wide range of green vegetables. There is therefore a risk that leafminer eggs or larvae (and potentially pupae and hitchhiking adults) could be spread to a new area via the movement of infested leafy vegetables. Mines do not occur on fruiting structures, with the exception that <i>Liriomyza huidobrensis</i> can form mines in the pods of snow peas and disrupt market access (Gitonga et al. 2010). Thus, movement of snow pea pods may also pose some risk.	Visual inspection for the presence of adults, larvae and pupae should be carried out. To reduce the potential risk of spread, plant material should not be moved from infested to non-infested areas without undertaking suitable treatments to reduce the risk of pest spread.

2.8.2 Predicting spread potential

Cesar Australia has created a model of the potential rate of area invasion by the serpentine leafminer at entry points across Australia, the results of which are summarised below (Figure 4). A copy of the relevant reports and/or published manuscripts are available from Hort Innovation or from Cesar Australia (<http://cesaraustralia.com/contact-us/>).




Figure 4 Rate of spread from locations across Australia. For each grid cell an incursion is simulated with the mean predicted area invaded by *L. sativae* in one year shown by the colour gradient. The mean predicted area is estimated from 10 replicated simulations. Incursions were commenced in July for a winter incursion and January for a summer incursion.

2.9 Risk of economic impact within Australia

Cesar Australia has created a model of the potential economic impacts of the serpentine leafminer in Australia, the results of which are summarised below (Table 4, Table 5, Figure 5). A copy of the relevant reports and/or published manuscripts are available from Hort Innovation or from Cesar Australia (<http://cesaraustralia.com/contact-us/>).

Table 4 Impact potential of serpentine leafminer on affected commodities in terms of international collated reports of proportions crop value lost due to direct feeding damage.

Host crop	proportion crop value lost			Sources
	Min	Max	Mean	
Beans	0.70	1.00	0.70	Rauf et al. 2000
Broccoli	0.10	1.00	0.10	Shepard et al. 1998
Capsicum	0.50	1.00	0.52	Spencer 1973; Elmore and Ranney 1954
Celery	0.15	1.00	0.46	Rauf et al. 2000; Weintraub et al 1998; Rauf et al. 2002
Cucumber	0.70	1.00	0.70	Rauf et al. 2000
Cut flowers	0.43	1.00	0.56	Rauf et al. 2000; Torres et al 1995

Potatoes	0.12	1.00	0.56	Rauf et al. 2000; Kwon et al. 2019; Waterhouse and Norris 1987; Chavez 1987; Chiluwal et al. 2012; Rauf et al. 2001; Shepard et al. 1998
Spinach	0.50	1.00	0.68	Rauf et al. 2000; Spencer 1973; Lange et al 1957
Tomatoes	0.00	1.00	0.15	Alves et al. 2017; Lopes et al 2019; Rauf et al. 2003

Table 5 Australia wide accumulated unmitigated impacts in millions of dollars after 3 years resulting from a spring (September) incursion of serpentine leafminer at key entry points across Australia. Simulations were replicated 10 times with means and standard deviations shown in parentheses. The proportion crop impact was fixed at 10% in order to explore variability due to incursion location and the size and distribution of different industries. Thus, host preferences of the pests do not influence the predicted crop impacts, and impacts of low preference hosts may be overestimated while impacts of high preference hosts may be underestimated. Predictions were made using ESIM framework and are the culmination of the establishment, spread and impact modules (with each level of risk building upon the previous). For impacts calculated at state levels, see Appendix 4, Table 20, for additional information. All cells with a value of 0.00 represent unmitigated impacts less than \$10,000.

Crop	Incursion Location					
	Bundaberg	Darwin	Devonport	Melbourne	Perth	Sydney
Beans	1.50 (0.00)	0.00 (0.00)	1.24 (0.17)	0.05 (0.02)	0.00 (0.00)	0.00 (0.00)
Broccoli	0.00 (0.00)	0.00 (0.00)	0.50 (0.21)	3.49 (1.63)	0.01 (0.01)	0.01 (0.03)
Brussels	0.00 (0.00)	0.00 (0.00)	0.33 (0.04)	0.03 (0.08)	0.00 (0.00)	0.00 (0.00)
Cabbages	0.00 (0.00)	0.00 (0.00)	0.12 (0.00)	0.34 (0.16)	0.00 (0.00)	0.02 (0.03)
Capsicum	1.69 (0.00)	0.00 (0.00)	1.50 (0.03)	0.01 (0.02)	0.00 (0.00)	0.00 (0.00)
Carrots	0.00 (0.01)	0.00 (0.00)	3.73 (0.03)	0.03 (0.03)	0.02 (0.03)	0.00 (0.01)
Cauliflowers	0.00 (0.00)	0.00 (0.00)	0.97 (0.00)	2.00 (0.96)	0.00 (0.01)	0.01 (0.01)
Flowers	0.89 (0.07)	0.00 (0.00)	1.40 (0.85)	0.47 (0.51)	0.01 (0.00)	0.45 (0.47)
Lettuces	0.07 (0.00)	0.00 (0.00)	0.27 (0.81)	2.00 (0.91)	0.02 (0.02)	0.05 (0.05)
Melons	3.98 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Nurseries	0.99 (0.18)	0.00 (0.00)	1.60 (0.66)	1.95 (1.31)	0.03 (0.01)	0.76 (0.56)
Onions	0.00 (0.00)	0.00 (0.00)	3.72 (0.01)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Peas	0.29 (0.00)	0.00 (0.00)	0.68 (0.14)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Potatoes	2.40 (0.00)	0.00 (0.00)	6.02 (0.08)	0.30 (0.37)	0.00 (0.01)	0.01 (0.02)
Pumpkins	0.35 (0.00)	0.00 (0.00)	0.14 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Tomatoes	5.50 (0.00)	0.00 (0.00)	0.35 (0.02)	0.00 (0.01)	0.00 (0.00)	0.02 (0.02)
Total	17.68 (0.25)	0.00 (0.00)	22.57 (1.81)	10.67 (4.44)	0.09 (0.08)	1.35 (1.00)




Figure 5 State level accumulating unmitigated impacts in millions of dollars across all host crops after 3 years resulting from a spring (September) incursion of serpentine leafminer at key entry and establishment points. For state level impact accumulation by crop type, see Appendix 4, Figure 9 for additional information

3. DIAGNOSTIC INFORMATION

Morphological diagnosis to a species level requires adult male flies. These can be collected by sweep netting, yellow sticky traps or collecting leaves with larvae in them and allowing the larvae to develop into adults. Alternatively, molecular tools can be used to identify larvae and adult flies.

A National Diagnostic Protocol for serpentine leafminer does not currently exist. However, an International Plant Protection Convention (IPPC) Diagnostic Protocol (IPPC 2016) and a European and Mediterranean Plant Protection Organization (EPPO) diagnostic protocol for *Liriomyza* spp. (EPPO 2005) are available. These diagnostic protocols should be referred to for the diagnosis of suspected *Liriomyza* spp. It should be noted that these protocols do not include information on environmental DNA (eDNA). Additional information on the use of eDNA for *Liriomyza* spp. leafminer diagnosis can be found at Pirtle et al., *in press*.

An information portal for polyphagous agromyzid leafminer identification is available at <https://keys.lucidcentral.org/keys/v3/leafminers/index.htm>, which includes species-specific fact sheets, and a pictorial Lucid3 key that allows users to make a preliminary identification of pest agromyzids and to be able to distinguish these species from non-pest endemic agromyzids. A factsheet for *Liriomyza huidobrensis* is accessible here:

https://keys.lucidcentral.org/keys/v3/leafminers/key/Polyphagous%20Agromyzid%20Leafminers/Media/Html/Liriomyza_huidobrensis.htm

3.1 Diagnostic considerations

Considerations when triaging samples of suspected serpentine leafminer for diagnostics include:

- One of the greatest challenges in detection and subsequent management of *Liriomyza* species overseas has been the difficulty in correctly identifying the species (Reitz et al., 2013), which can have considerable overlap in adult morphology, host range and damage symptoms left on a plant. In the presence of indigenous leafminer, it may be difficult to detect a new incursion, or determine the limits of an infestation (Powell 1981).
- There are also several difficult to distinguish Agromyzid species present within Australia that are morphologically indistinguishable from serpentine leafminer to a non-expert (including *L. brassicae* and *L. chenopodii*). This is also true for some Agromyzids that are exotic to Australia. In many cases, specimens must be distinguished by expert taxonomists, or via molecular means.
- Within Australia, there are at least 50 leafminer species (family Agromyzidae) (and presumably many more that have not yet been described) that create leaf mines that look similar to serpentine leafminer and may appear on the same host plants. Some of the most common include *Liriomyza brassicae* (which shares several brassica hosts with serpentine leafminer), *Liriomyza chenopodii* (which shares cultivated beets with serpentine leafminer), and *Phytomyza syngenesiae* (which shares cultivated asters with serpentine leafminer, but the adult form is readily distinguishable from serpentine leafminer) (Appendix 2). Moreover, there are an unknown number of additional Dipteran and Lepidopteran (Figure 8) species that share the leaf mining habit.
- Of these species that create similar leaf mining damage, some can be distinguished from serpentine leafminer via their mine characteristics, and their pupal or adult forms. (see Appendix 2, IPCC 2016, Salvo and Valladares 2004). However, the high degree of individual variation between each leaf mine means there can always be exceptions to these guidelines. For example, while serpentine leafminer pupate outside the leaf, there are rare instances where a larvae will fail to emerge and complete pupation within the leaf mine.
- Due to the high degree of morphological overlap of adults and visual overlap of damage, sample collection (of flies of any life stage and/or leaf mines) will be paramount for identifying and delimiting potential serpentine leafminer incursions.

- Surveyors should be familiar with the serpentine leafminer host plant list (Appendix 1), as well as with commonly observed signs of indigenous leafminer damage and the extent of known overlap with indigenous Agromyzids for common host genera (Appendix 2) before conducting serpentine leafminer surveillance. For example, damage caused by the closely related *L. brassicae* is essentially indistinguishable from serpentine leafminer and is very common on brassica crops (Appendix 2). Crops such as tomato, cucumbers and melons, on the other hand, have little to no reported overlap with indigenous Agromyzids (Appendix 2). Mines detected on these crops during surveillance activities should be considered highly suspicious.

4. SURVEILLANCE AND COLLECTION OF SAMPLES

4.1 Surveillance

4.1.1 Technical information for planning surveys

Detection and delimiting surveys are required to determine the extent of outbreaks, ensure areas free of the pest retain market access, and ensure appropriate quarantine zones are established to contain outbreaks.

Initial surveillance priorities for serpentine leafminer should consider the following:

- Visual surveillance for the leaf mines created by serpentine leafminer is expected to be the most reliable way to detect new incursions. This is because the mines created by leafminer larvae persist permanently on a leaf, extending the detection window beyond the lifespan of the fly itself. In comparison, sweep netting requires active populations be found, as do yellow sticky traps. While sticky traps can be left in place for extended periods to increase the chance of catching an adult, in open field settings, relying solely on sticky traps would be impractical due to the very large number of traps that would be needed to be placed and processed, to reach a statistically acceptable detection likelihood for small or isolated populations.
- In closed cropping systems, sticky traps become much more practical, but should still be accompanied by visual surveillance.
- Sticky traps will have a high level of bycatch (i.e. other insects will be caught in the trap), which can increase processing time for manual identifications. However, molecular diagnostic methods are also available for bulk invertebrate samples collected via a sticky trap (see Section 4.2.2.2).
- When undertaking visual surveillance in plant crops, the leaf mines are the most conspicuous symptoms to look for, though stippling from oviposition and feeding may also be apparent.
- Mines are expected to be more common on the edges of fields and in the upper portions of the plant (Weintraub 2001).
- The serpentine leafminer has an extensive list of weed hosts (see Section 2.3) and therefore surveillance in broadleaf weeds as well as cultivated crops should be conducted.
- Larvae, pupae and adults can be identified using molecular methods (refer to Section 3).
- Only adult male leafminers can be identified using morphological methods.
 - Adults can be captured using yellow sticky traps, sweep netting (IPPC 2016), or rearing from infested material. The latter is the most practical and reliable method for collecting adult flies. Infested leaves can be placed into an insectarium to develop into adults (IPPC 2016). Techniques for rearing agromyzids are described in Griffiths (1962) and Fisher et al. (2005).
- Collectors should be made aware of common signs of leafminers and note symptomatic plants when conducting surveillance for serpentine leafminer.
- Australia is already home to several leaf miner species, both native and naturalized, which can be confused for serpentine leafminer. Surveyors should be made aware of the native and

naturalized species they might expect to see in the particular region and crops they are surveying.

- Early detection surveillance will be most effective if timed when serpentine leafminer population growth potential is greatest, such as during warm (around 20-30°C) conditions. The seasonality of serpentine leafminer activity at major centers is provided in Table 6.

Table 6 Major Australian growing regions showing risk seasonality for serpentine leafminer activity across the year. Green means highest establishment risk (where the population growth potential was greater than 50% of the maximum predicted growth rate) yellow means lower establishment risk (where the population growth potential was between 1 and 50% of the maximum), and red means lowest establishment risk (where predicted growth rates are negative, meaning populations are predicted to decline).

State	Incursion point	Seasonality											
		Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Western Australia	Perth	Red	Red	Red	Red	Yellow	Green	Yellow	Green	Green	Red	Red	Red
	Geraldton	Red	Red	Red	Red	Red	Green	Green	Green	Green	Red	Red	Red
	Carnarvon	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red
	Bunbury	Red	Red	Red	Green	Red	Green	Yellow	Yellow	Green	Red	Red	Red
	Albany	Red	Green	Green	Green	Yellow	Yellow	Yellow	Yellow	Green	Green	Yellow	Yellow
Victoria	Thorpdale	Green	Green	Green	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Green	Green	Green
	Melbourne	Yellow	Yellow	Red	Green	Yellow	Yellow	Yellow	Yellow	Green	Red	Red	Green
	Ballarat	Green	Yellow	Red	Green	Yellow	Yellow	Yellow	Yellow	Green	Red	Red	Green
Tasmania	Smithton	Yellow	Yellow	Green	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Green	Green	Green
	Devonport	Green	Green	Green	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Green	Green	Green
South Australia	Mt. Gambier	Green	Green	Red	Green	Yellow	Yellow	Yellow	Yellow	Yellow	Green	Green	Green
	Loxton	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red
	Adelaide	Red	Red	Red	Red	Red	Red	Red	Red	Red	Yellow	Red	Red
Queensland	Toowoomba	Red	Red	Yellow	Green	Green	Green	Yellow	Green	Red	Red	Green	Yellow
	Bundaberg	Red	Red	Red	Green	Green	Green	Green	Green	Red	Red	Green	Yellow
	Bowen	Red	Red	Red	Green	Green	Green	Yellow	Red	Red	Red	Red	Red
	Atherton	Red	Green	Green	Green	Green	Green	Yellow	Yellow	Yellow	Red	Red	Red
Northern Territory	Katherine	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red
	Darwin	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red
New South Wales	West Sydney	Red	Red	Green	Yellow	Yellow	Yellow	Yellow	Yellow	Green	Green	Yellow	Yellow
	Sydney	Red	Red	Green	Green	Green	Yellow	Yellow	Yellow	Yellow	Red	Red	Red
	Griffith	Red	Red	Red	Green	Green	Yellow	Yellow	Yellow	Green	Red	Red	Red

4.1.1.1 Surveillance tools

Yellow sticky traps and sweep netting can be used as supplementary methods to visual searches during delimiting surveillance activities, particularly when suspicious damage has been observed, or for monitoring existing outbreaks.

- Sweep netting is expected to be most effective when employed near recently active leaf mines, and is less effective near old leaf mines (which tend to be darker coloured than fresh mines, see Figure 6). Windy conditions can also reduce the effectiveness of sweep netting as adult leafminers tend to remain low in foliage during windy conditions (Foster 1986).



Figure 6 Younger mines (on the left) are more white-coloured with more distinct black trails of frass. Frass becomes less distinct as mines age, and mines develop a brown colour (right). Younger mines are expected to give the best sweep netting success. Images: Elia Pirtle, Cesar Australia Pty Ltd

- Agromyzid flies are attracted to the colour yellow (Chavez et al., 1987), and can therefore be captured on yellow sticky traps. Yellow sticky traps are more effective for *Liriomyza* adults than other types of traps, such as funnel traps and yellow water pans (Chavez and Raman 1987). Vacuum sampling has also been shown to be a less effective means of sampling adult leafminer (Weintraub 2001).
- A great deal of effort overseas has been dedicated to improving the effectiveness of yellow sticky traps for *Liriomyza* adults (Heinz et al 1992) including the modifications of size, shape, adhesives, lures, height and orientation. For example, several studies report a strong effect of trap height on the number and species trapped (Trumble and Nakakihara 1983, Zehnder and Trumble 1984, Zoebisch and Schuster 1990), however these results do not always appear consistent and may be difficult to extrapolate across different crop types. Moreover, optimal height may vary considerably between *Liriomyza* species (Zehnder and Trumble 1984).
- Some considerations when using yellow sticky traps for *Liriomyza* surveillance are provided below:
 - Traps should be placed (vertically oriented) at the edges of paddocks.
 - Optimal trap height may vary with crop type. Studies on potato have found that traps placed at the crop's canopy height captured more flies than traps placed near the ground or above the crop's canopy (Weintraub and Horowitz 1996).
 - Traps should be left for a minimum of 24 hours, to ensure traps are in place during the peak flight time for serpentine leafminer, which occurs just after sunrise (Weintraub and Horowitz 1996).
 - Traps may be left in place for up to one month, at which point they should be collected, to prevent excessive degradation of DNA and allow for molecular confirmation of any leafminers caught on the trap. If traps are placed outdoors and exposed to the elements and/or high levels of bycatch they should be collected sooner than one month.

- Experimental lures developed from the extracted volatiles of known plant hosts have been shown to be attractive to *Liriomyza*. For example, lures made from spruce, basil, juniper or clove oil have been shown to attract serpentine leafminer (Gorski 2005).

4.1.2 Delimiting surveys in the event of an incursion into new areas

Following a new detection of serpentine leafminer, the following guidelines are recommended for delimiting surveillance:

- Surveillance to delimit a detection of serpentine leafminer should take into account tracing information as outlined in Section 5.1 to determine potential pathways for movement of material from the site of the initial detection.
- Surveillance should be a combination of the following:
 - Visual inspection in high risk areas (e.g. areas with suitable hosts).
 - Sweep nets to collect adults around infested host plants.
 - Yellow sticky traps, particularly in closed cropping systems and around infested plants/crops.
- Surveillance should be accompanied with awareness material, signs and personal visits to households and businesses within the surveillance zone and any buffer zones.

The suggested surveillance procedure for serpentine leafminer is as follows:

1. Delimiting surveillance should be conducted immediately upon detection of any incursions within a radius within 1 km of the incursion site, based on natural spread dynamics of agromyzid leafminer. The Surveillance zone may be increased as deemed necessary for the individual response.
2. At each site, choose a crop that is a preferred host plant of serpentine leafminer (refer to Section 2.3).
3. Choose a survey path that preferences the following areas if possible:
 - Paddocks and weeds near transport routes and unloading areas. These are preferred, because a major long-distance pathway of leafminer spread is human assisted movement.
 - Paddock edges should be included in the survey path, with the 'incoming wind side' preferred, since short distance movements of leafminer can be driven by wind.
 - Broadleaf weeds along paddocks should be included in the survey path.
4. Leafminer damage may be patchy at low densities.
5. At low densities, leaf mines have a relatively low detectability, despite their characteristic appearance. Slower search speeds will improve detection likelihood, however, when time is limited, search speed trades off with the total area covered (reducing the number of possible encounters). This trade-off is important in realistic paddock monitoring scenarios.
 - When time is not limited: To achieve a detection probability of 20%, 50%, and 80% respectively, when leaf mine density is about 1% of plants affected, the mean (and 95% CI) search time required per metre is 11.2 (4.8, 20.5), 34.8 (15.0, 63.8), and 80.8 (34.9, 148.2) seconds spend 1 minute searching per 1 m² of area (based on data collected during vegetable leafminer survey effort trials).

- When time is limited: The optimal search speed, balancing area coverage with sensitivity, when leaf mine density is about 1% of plants, is about 10 seconds per 1 m² of area. Further benefits are not achieved by increasing search speed beyond this level to achieve greater area coverage. (These recommendations are based on data collected during Vegetable leafminer (*L. sativae*) survey effort trials – other *Liriomyza* leafminers are expected to require similar effort)).
6. A survey plan should take into consideration any variables regarding the environment or surveyor that might reduce detectability and increase the effort spent accordingly. These variables might include:
 - *Plant type*
 - i. *Leaf density*: leafminer damage may be harder to spot in plants that are very dense and leafy, than in plants with larger, sparser or flatter leaves.
 - ii. *Plant age*: it may be easier to spot leaf mines in young plants with few leaves and more open space between plants than in larger mature, overlapping plants.
 - iii. *Colour*: leafminer damage may be harder to spot against light coloured leaves than against darker leaves.
 - iv. *Height*: Leafminer damage in plants that are at eye level may be easier to spot than crops that are low to the ground.
 - v. *Time of day/direction of survey relative to the sun*: Survey success may be higher when surveyors are walking with their back to the sun, rather than walking into the sun.
 - *Residues on plants*: Chemical residue on plants may reduce the contrast between the mine and the leaf, or obscure mine patterns with background residue patterns.
 - *Damage from other pests/disease on plants*: damage caused by other pests or disease may obscure leaf mines, or lead to false positive detections.
 - *Surveyor experience*: Inexperienced surveyors may have a learning curve.
 - *Weather conditions*: Surveys during poor weather should be avoided when possible.
 7. If suspicious damage is detected, samples should be collected (see Section 4.2), and yellow sticky traps should be placed around the affected area, in an attempt to capture adults and diagnose the fly responsible for the damage.
 - a. If serpentine leafminer are confirmed, visual surveillance along with sticky traps should be used to monitor the edges of the outbreak area.
 8. If no suspicious damage is detected, the initial round of delimiting surveillance should be followed with a second round occurring after one week.
 - a. This second round is necessary as leafminer eggs are undetectable in the leaf, and hatch after about 2-5 days. A crop that appears clean during the initial survey may still develop symptoms within a week.
 - b. After the first revisit, any further revisits should occur on a monthly basis. One lifecycle of the serpentine leafminer is estimated to take approximately 2 to 4 weeks, dependent on host and temperature.

4.1.3 Activities for general surveillance immediately following a detection

Given the wide host range of this pest and its noticeable impact it may be a suitable target for general surveillance programs that request submission of images and/or samples from the public.

To establish a general surveillance program, the following will be required:

- Use of online or app reporting tools such as MyPestGuide to promote the submission of reports of suspected detections.
- Factsheets to provide information on the pest, symptoms, impacts and reporting mechanisms.
- Media releases to describe the impact of the pest, surveillance programs and activities within the response program.
- Information for industry communication channels including articles for industry newsletters, magazines and websites, information for Twitter feeds and Facebook, and presentations for industry talks.
- A website to provide information for the public and for commercial businesses and links to other relevant sites.
- Release and promotion of information on details for physical sample collection and submission, as well as information on how to take and submit images of flies/leaf mines.
- Broader awareness campaigns should consider literature (brochures and factsheets) in several languages.
- A non-expert friendly surveillance and sample collection guide for vegetable leafminer, which also has relevance for the serpentine leafminer, as well as other useful information, can be found [here](#).

4.2 Collection and treatment of samples

4.2.1 If collecting for morphological diagnosis

Of the four life stages (egg, larva, pupae and adult) only adult males are identifiable to a species level using morphological features.

4.2.1.1 Collection of adult male specimens

Adult flies can be collected with a vacuum sampler or sweep net. Adult flies are normally found on the foliage. However, the most practical and reliable method is the collection of leaves with mines containing mature larvae or pupae which can then be reared until adult flies emerge (refer to Section 4.2.1.3).

4.2.1.2 How to collect sticky traps for identification

Sticky traps should be left in the field for up to one month (but preferably 2 weeks or less), to allow morphological identifications. Upon collection, sticky traps should be folded lightly in half sticky side inwards (not pressed shut), and placed in a sealed, padded plastic bag. Traps should be sent within a week of collection (and kept in cool, dry place in the meantime, but not frozen).

4.2.1.3 How to collect plant samples for rearing

For rearing of adult flies, mined leaves containing pupae or mature larvae can be collected into an

insectarium and kept in a constant temperature room for regular checking. Simple insectariums include sealed partially inflated ziplock bags, or sealed rigid plastic containers or jars. A piece of paper towel should be placed into the insectarium along with the infested leaves to regulate humidity. Once pupae emerge from the leaves, they should be transferred carefully with a wet paintbrush to a new rigid container, which contains a layer of damp sand but no leaf material. Pupae can be placed directly onto the damp sand. Techniques for rearing agromyzids are described in the IPCC diagnostic protocol for *Liriomyza* (IPPC 2016, available from [here](#)).

4.2.1.4 How to preserve plant samples for identification

Identifying the plant species affected by suspected serpentine leafminer damage can aid diagnoses, particularly when surveying in weeds. To preserve a plant for identification, leaves can be stored between sheets of dry newspaper. However, if plant material with suspected leaf mines is pressed and dried, there will be no chance of rearing out adult flies from those leaf mines. Therefore, whenever plant identity is in question, infested material should be stored in an insectarium for rearing, and an accompanying pressed sample of uninfested leaves should be taken to facilitate for identification of the host.

4.2.1.5 How to preserve leafminers

Adults caught in sweep nets or reared in insectariums can be placed in 70% ethanol and stored indefinitely for morphological purposes. For adults reared in insectariums, it is advised to collect these adults in the afternoon, as emergences tend to occur in the morning, and enough time should be given for wings to fully expand before collection.

4.2.1.6 How to transport leafminers

Vials of ethanol should be sealed to avoid leakage and packed with cushioning material in a strong box.

For information on what can or cannot be sent by post refer to the Australia Post website:

https://auspost.com.au/content/dam/auspost_corp/media/documents/dangerous-and-prohibited-goods-guide.pdf.

4.2.1.7 How to transport plant samples

Plant samples may be collected if the host needs to be identified. Queensland Herbarium (2016) provides information on how to collect and preserve herbarium specimens. Samples should then be taken to state herbaria for identification.

For information on what can or cannot be sent by post refer to the Australia Post website:

https://auspost.com.au/content/dam/auspost_corp/media/documents/dangerous-and-prohibited-goods-guide.pdf.

4.2.2 If collecting for molecular diagnosis

All life stages can potentially be diagnosed using molecular methods, including empty leaf mines.

4.2.2.1 Collection of specimens

Adult flies can be collected with a vacuum sampler or sweep net. Adult flies are normally found on the

foliage. However, as other life stages can be diagnosed using molecular methods leaves with mines containing pupae or mature larvae should also be collected for analysis. Rearing these life stages in an insectarium is not necessary.

4.2.2.2 How to collect sticky traps

Upon collection, sticky traps should be either:

- Placed inside a plastic sleeve that has been cut along one long side and along the bottom, the trap and sleeve can then be sealed in a plastic zip lock bag.
- Folded in half, sticky side inwards, and placed in a sealed plastic bag.

Traps should be sent within a week of collection (and kept in a cool, dry place in the meantime, but never frozen).

4.2.2.3 How to collect plant samples

Plant samples with suspected serpentine leafminer damage should be collected for molecular confirmation. If it is possible to discern a live larvae inside a leaf mine (which can be spotted through the leaf mine as a small yellow maggot, either by eye or with a hand lens, these samples are always preferable (which can be spotted through the leaf mine as a small yellow maggot, either by eye or with a hand lens, see Figure 7). However, mined leaves that appear empty can also be collected. Empty mine samples that look youngest should be preferred, as they have the greatest chance of containing a live larva (see Figure 6).



Figure 7 A larva inside a leaf mine. Larvae are visible to the eye in later stages of development. Images: Elia Pirtle, Cesar Australia Pty Ltd

4.2.2.4 How to preserve plant samples

Mined leaves containing pupae or mature larvae, or mined leaves that appear empty can all be placed straight into 100% Ethanol. Small mined leaves should be rolled and placed into the tube of absolute ethanol. For larger leaves, the mine should be cut out, with care taken to avoid cutting across the mine, and the cut out should be rolled and placed into the tube of absolute ethanol. These samples should be frozen and mailed within a week of collection for molecular diagnosis.

Due to the very small amounts of DNA within leaf mines, the odds of sample contamination is much higher than when working with whole flies. Careful hygiene must be employed when collecting and

preserving suspicious leaf mine samples. Mines from different host plants or different locations should never be stored in the same container. Samples should either be immediately preserved into on site, or should be placed into unused, clean, sealed plastic bags on site and preserved within the day. If samples from multiple hosts or sites are being processed together at a field lab location, all equipment must be sterilised between each host/location (i.e. cleaning counter tops, sterilizing tweezers and other tools).

Leaf mines may also be preserved onto FTA cards, however DNA quality is somewhat lower than for preservation into 100% ethanol (Pirtle et al *in press*). To preserve leaf mines onto FTA cards, remove the mined leaf from plant and push the leaf mine face down onto the FTA card. Smear the mine thoroughly on the FTA card, for about 30 seconds, and then close the card and place in small sealed plastic bag. Do not rub mines from different host plants/sites onto the same FTA card, and do not store FTA cards from different host plants/sites in the same sealed bag. Store the FTA cards in a cool, dark, dry place and send for analysis within a week of collection

4.2.2.5 How to preserve leafminers

Specimens required for molecular diagnostic work, including larvae, pupae or adults, should be killed and preserved in absolute ethanol, frozen at -80°C, or preserved on an FTA card.

4.2.2.6 How to transport leafminers and plant samples

Vials of ethanol should be sealed to avoid leakage and packed with cushioning material in a strong box.

For information on what can or cannot be sent by post refer to the Australia Post website:

https://auspost.com.au/content/dam/auspost_corp/media/documents/dangerous-and-prohibited-goods-guide.pdf.

5. IMMEDIATE RESPONSE TO DETECTION

5.1 Tracing

Detection and delimiting surveys are required to delimit the extent of the outbreak, support areas free of the pest to retain market access, and ensure that appropriate quarantine zones are established.

Extensive tracing (trace forward and trace back) may be feasible as *Liriomyza* spp. can be readily dispersed by the movement of infested plant material, soil and crop debris. The focus should be on high risk linkages including premises linked directly with the initial detection, particularly where movements of plant material (such as nursery plants and seedlings), or soil have occurred as these pathways may move serpentine leafminers over long distances.

Further information on possible risk pathways are presented in Section 2.8 and in Table 2.

5.2 Quarantine and movement controls

If Restricted or Quarantine Areas are practical, no plant material should be moved from the infested to unaffected areas without first being inspected and appropriately treated. The size of the Restricted or Quarantine Area will be dependent on the type and scale of the incursion. However, a zone 1 km in radius should be considered for initial delimiting surveillance activities (Section 4.3). This zone can then be expanded as needed depending on the specific situation.

Voluntary movement control should be considered for urban/residential detections. Voluntary controls would involve negotiation with residents to undertake inspection and treatment of goods prior to movement from infested areas. Residents should be advised on measures to minimise the inadvertent transport of the pest from the infested area to unaffected areas. Voluntary compliance is likely to be implemented for urban areas using awareness campaigns to highlight high risk goods/situations and appropriate treatments.

5.3 Destruction strategy

5.3.1 Priorities and considerations

If eradication of an outbreak is considered feasible, the priority will involve the destruction of the pest. Adult leafminers may be present around host plants; eggs, larvae and pupae may be present in the leaves of host plants, and pupae may be present within the first 5 cm of soil under host plants.

Considerations include:

- The infested plants may need to be sprayed with a suitable insecticide (refer to Section 5.3.3), then bagged and then buried (or tilled/ploughed directly into the soil if a large area is affected).
- Other host plants in the immediate area should be inspected and treated as needed.
- Soil and plant media in greenhouses should be treated (heat or chemically) to destroy any pupae that may be present in the soil.

- Machinery and equipment used on site may need to be washed down to minimise the potential risk of spreading the pest between sites.

5.3.2 Decontamination of tools and equipment

Machinery, equipment, tools and clothes should be cleaned and disinfected between sites to minimise the potential to spread leafminers, and other pests and diseases that may be present on the site, to new areas.

For points to consider when decontaminating, refer to the PLANTPLAN Guideline document: Disinfection and Decontamination (PHA, 2015). which includes guidelines for the destruction and decontamination of tools used when working with emergency plant pests.

Several state agriculture departments have also produced publications on cleaning down machinery and vehicles and should be referred to if decontamination is required, examples include:

- www.dpi.nsw.gov.au/_data/assets/pdf_file/0010/545554/procedure-decontamination-vehicles-and-equipment.pdf.
- www.daf.qld.gov.au/_data/assets/pdf_file/0011/58178/cleandown-procedures.pdf.
- <http://agriculture.vic.gov.au/agriculture/pests-diseases-and-weeds/weeds/weedstop-vehicle-hygiene-program/machinery-hygiene>.

5.3.3 Chemical control

Overseas *Liriomyza* leafminers are most damaging when parasitoids are removed due to the use of non-selective insecticides (Cesar Australia 2018; Murphy and LaSalle 1999). For the ongoing management of leafminers it will therefore be important that chemicals are selected that are reasonably selective and have low toxicity towards key parasitoids.

The European and Mediterranean Plant Protection Organization (EPPO) provides guidance (EPPO 2009) on how to manage and eradicate Vegetable leafminer infestations in Europe, which are broadly applicable to other *Liriomyza* spp., such as the serpentine leafminer. For Vegetable leafminer, the EPPO recommends that foliar sprays are applied at weekly intervals until control has been achieved. Importantly the EPPO suggests rotating chemicals and modes of action (where possible) each treatment (Table 7). In greenhouse environments the use of a suitable space treatment is also suggested (EPPO 2009). Based on this it would be ideal if pesticides from different mode of action groups are available for each host crop, as this would allow pesticides to be rotated rather than relying on a single product being continuously used to manage/treat the pest.

Table 7 An ideal chemical rotation schedule used for eradication of *Liriomyza* leafminers in Europe (refer to EPPO 2009)

WEEK	PESTICIDE USED TO TREAT INFESTED AREA
Week 1	Chemical A
Week 2	Chemical B
Week 3	Chemical C
Week 4 onwards	Repeat above sequence

Pesticides registered for the control of serpentine leafminer overseas include the following actives:

- Abamectin
- Azadirachtin
- Chlorantraniliprole
- Cyantraniliprole
- Cyromazine
- Diazinon
- Emamectin benzoate
- Indoxacarb
- Permethrin
- Spinetoram
- Spinosad
- Spirotetramat

It should be noted that Australian Pesticides and Veterinary Medicines Authority (APVMA) permits will need to be put in place before these pesticides can be used for the control of serpentine leafminers. Permits are being applied for as part of Horticulture Innovation Australia Project MT16004 (RD&E program for control, eradication and preparedness for Vegetable leafminer), and these are noted in Table 8.

As the serpentine leafminer can affect many plant species (including agricultural and ornamental species) (refer to Section 2.3), it is important that all potential hosts are covered by at least one, but preferably multiple, pesticide permits. Appendix 3 provides a summary of the pesticides that can be used on each crop type and host and should be used as a guide when treating new serpentine leafminer outbreaks, or if eradication is unsuccessful, managing the pest. As new permits are approved, they will be made publicly available on the APVMA permit database:

<https://portal.apvma.gov.au/permits>.

Table 8 Insecticide permits that have been applied for or granted by the APVMA for the control of serpentine leafminer in Australia (as of July 2020)

CHEMICAL	CHEMICAL ACTIVITY	LIFECYCLE STAGES CONTROLLED	CROPS COVERED ¹	PROPOSED RATE	PERMIT NUMBER	EXPIRATION DATE
Cyromazine (MOA 17)	Contact and translaminar activity	Not effective against adults but being an insect growth regulator is very effective against larval stages (Cesar Australia 2018b). Experiments on the related <i>L. trifolii</i> found that Cyromazine was very effective against first and second instar but less effective against late instar larvae (Yathom et al., 1986).	Nursery stock (non-food)	333 g in 20 L water/10 tonne potting mix	PER83506	31 October 2022
			Fruit trees (non-bearing)	950 g/ha or 95 g/100 L applied in 1,000 L/ha.		
			Broccoli	1 kg/ha or 150g/100L water	PER81867	30 November 2023
			Fruiting vegetables – cucurbits			
			Fruiting vegetables – other than cucurbits (excluding mushrooms and corn)			
			Head lettuce			
			Legume vegetables			
			Root and tuber vegetables			
			Stalk and stem vegetables			
			Brassicas (excluding broccoli)			
			Leafy vegetables (excluding head lettuce)			
			Note: this crop must be destroyed if treated and must not be made available for human consumption			
			Note: this crop must be destroyed if			

¹ Refer to Appendix 3 for detailed information on chemical control options.

CHEMICAL	CHEMICAL ACTIVITY	LIFECYCLE STAGES CONTROLLED	CROPS COVERED ¹	PROPOSED RATE	PERMIT NUMBER	EXPIRATION DATE
			treated and must not be made available for human consumption			
Dimethoate (MOA 1B)	Contact and systemic activity	Dimethoate is reported to control <i>Liriomyza</i> spp. adults and larvae however resistance is known overseas (Tran and Takagi 2005)	Pulse crop (grains)	75ml (400 g a.i/L products)/100L water	PER89184	31/3/2025
Eamectin benzoate (MOA 6)	Contact and translaminar activity	Same mode of action as abamectin. Reported efficacy against larvae, adults and eggs (Cesar Australia 2018b)	Brassica vegetables (including broccoli, Brussels sprouts, cabbage, cauliflower)	250 to 300 g/ha (44 g a.i/kg products); or 650 to 780 mL/h (17 g a.i/L products) (suppression only)	PER87563	30 June 2024
Spinetoram (MOA 5)	Contact, systemic and translaminar activity	Spinetoram has shown efficacy against <i>L. sativae</i> adults and larvae (Shimokawatoko et al., 2012)	Vegetable legumes (peas and beans)	400 ml (48 g a.i)/ha	PER87878	28/2/2023
Spirotetramat (MOA 23)	Contact, systemic and translaminar activity	Affects the development of larvae and has little impact on adults (Ring 2019)	Capsicum	400ml (96 g a.i)/ha	PER88640	31/5/2023
			Celery			
			Chili			
			Eggplant			
			Green beans			
			Lettuce (head and leafy)			
			Parsley			
			Rhubarb			
			Snow & sugar snap peas			

CHEMICAL	CHEMICAL ACTIVITY	LIFECYCLE STAGES CONTROLLED	CROPS COVERED ¹	PROPOSED RATE	PERMIT NUMBER	EXPIRATION DATE
			Tomato			

Research undertaken as part of the Hort Innovation project MT16004 has identified the suitability of the following as potential control options for *Liriomyza* spp.:

- Azadirachtin (a systemic insecticide that acts as an insect growth regulator, MOA unknown) on ornamental and floriculture plants.
- Abamectin (an insecticide with contact and translaminar activity, MOA 6) on fruiting vegetables, cucurbits, leafy vegetables, legume vegetables, bulb vegetables, cabbage, celery, and rhubarb.
- Cyantraniliprole (a systemic insecticide, MOA 28) on bulb vegetables, fruiting vegetables, cucurbits, melons, and potatoes (note residue data will be required to support this use pattern).
- Diazinon (a contact insecticide, MOA 1B) on beans, melons, beetroot, lettuce, parsnips, tomatoes, and potatoes. Note can have significant impact on beneficial insects meaning it would likely be more suited to use in eradication programs than ongoing management of the pest.
- Dimethoate (a systemic insecticide, MOA 1B) on green beans, peas, tomatoes, capsicums, and potatoes. Note can have significant impact on beneficial insects and not fit well with IPM systems.
- Emamectin Benzoate (an insecticide with contact and translaminar activity, MOA 6) on brassicas, leafy vegetables, fruiting vegetables, cucurbits, and melons.
- Indoxocarb (an insecticide with contact and some translaminar activity, MOA 22A) on tomatoes.
- Permethrin (an insecticide with contact activity, MOA 3A) on tomatoes. Note can have significant impact on beneficial insects meaning it would likely be more suited to use in eradication programs than ongoing management of the pest.
- Pyrethrins (an insecticide with contact activity but very limited residual activity, MOA 3A) on brassicas, leafy vegetables, lettuce, fruiting vegetables, and legume vegetables.
- Spinetoram (an insecticide with contact, systemic and translaminar activity, MOA 5) on cotton.

Permits for these use patterns have not been applied to date. However, in the event of an incursion, emergency permits for these use patterns could be developed if required.

5.3.4 Physical control options

Destruction of host plants and deep ploughing of crop residues can assist with leafminer control as adult leafminers have difficulty emerging from pupae buried deeply in soil.

Removal of weed hosts of the leafminer and planting susceptible crops away from weed hosts can also assist with the ongoing management of the pest by removing/avoiding potential leafminer reservoirs.

5.3.5 Biological control

Although not specifically for eradication, biological controls will assist in lowering leafminer populations and will be an important part of the management of serpentine leafminers should eradication not be possible in the event of an outbreak.

Parasitoids often provide effective suppression of leafminers in the field when non-selective insecticides are not used (Cesar Australia 2018). History has also shown that invading *Liriomyza* populations are rapidly exploited by endemic agromyzid parasitoids (Cesar Australia 2018). For these reasons it will be important that any pesticides used for the ongoing management of leafminers are compatible with Integrated Pest Management (IPM) systems.

Weintraub et al., (2017) identified more than 100 species of parasitoids that are reported to affect serpentine leafminer (Table 9) these represent 7 families of organisms. Agromyzid parasitoids tend not to be very host-specific and Australia already has several endemic species that would likely be effective against the serpentine leafminer. *Diglyphus isaea* is an example of a parasitoid that occurs in Australia and is mass reared overseas for the biological control of *Liriomyza* species including serpentine leafminers.

The importation of biological control agents is not warranted at this time as the adventive parasitoids such as *D. isaea*, already occur in Australia and would likely be suitable for the management of leafminers assuming pesticides are selected appropriately and are compatible with IPM systems.

Table 9 parasites and parasitoids of serpentine leafminer noted in Weintraub et al., (2017)

PARASITE NAME	PARASITE NAME	PARASITE NAME	PARASITE NAME
<i>Agrostocynips clavatus</i>	<i>Dacnusa sibirica</i>	<i>Halticoptera helioponi</i>	<i>Opius</i> sp.
<i>Alloxysta</i> sp.	<i>Dacnusa</i> sp.	<i>Halticoptera patellana</i>	<i>Pediobius metallicus</i>
<i>Asecodes delucchi</i>	<i>Diaulinopsis arenaria</i>	<i>Halticoptera peviana</i>	<i>Phaedrotoma luteoclypealis</i>
<i>Asecodes</i> sp.	<i>Diaulinopsis callichroma</i>	<i>Halticoptera</i> sp.	<i>Phaedrotoma mesoclypealis</i>
<i>Bracon intercessor</i>	<i>Diaulinopsis</i> sp.	<i>Hemiptarsenus fulvicollis</i>	<i>Phaedrotoma scabriventris</i>
<i>Chrysocharis ainsliei</i>	<i>Diglyphus albiscapus</i>	<i>Hemiptarsenus ornatus</i>	<i>Phaedrotoma</i> sp.
<i>Chrysocharis bedius</i>	<i>Diglyphus begini</i>	<i>Hemiptarsenus</i> sp.	<i>Platynocheilus cuprifrons</i>

PARASITE NAME	PARASITE NAME	PARASITE NAME	PARASITE NAME
<i>Chrysocharis brethesi</i>	<i>Diglyphus crassinervis</i>	<i>Hemiptarsenus unguicellus</i>	<i>Phigalia incompletus</i>
<i>Chrysocharis c.f. aluta</i>	<i>Diglyphus intermedius</i>	<i>Hemiptarsenus varicornis</i>	<i>Phigalia katonis</i>
<i>Chrysocharis caribea</i>	<i>Diglyphus isaea</i>	<i>Hemiptarsenus zilahisebessi</i>	<i>Phigalia soemius</i>
<i>Chrysocharis flacilla</i>	<i>Diglyphus minoeus</i>	<i>Heteroschema</i> sp.	<i>Phigalia</i> sp.
<i>Chrysocharis ignota</i>	<i>Diglyphus pachyneurus</i>	<i>Moneucoela</i> sp.	<i>Proacrias</i> sp.
<i>Chrysocharis orbicularis</i>	<i>Diglyphus pedicellus</i>	<i>Neochrysocharis beasleyi</i>	<i>Proacrias thysanoides</i>
<i>Chrysocharis oscinidis</i>	<i>Diglyphus pedicellus</i>	<i>Neochrysocharis diastatae</i>	<i>Proacrias xenodice</i>
<i>Chrysocharis pentheus</i>	<i>Diglyphus poppoea</i>	<i>Neochrysocharis formosa</i>	<i>Quadrastichus liriomyzae</i>
<i>Chrysocharis pubicornis</i>	<i>Diglyphus pulchripes</i>	<i>Neochrysocharis okazakii</i>	<i>Quadrastichus</i> sp.
<i>Chrysocharis</i> sp.	<i>Diglyphus</i> sp.	<i>Neochrysocharis</i> sp.	<i>Sphegigaster</i> sp.
<i>Chrysocharis tristis</i>	<i>Diglyphus</i> sp. (near <i>intermedius</i>)	<i>Notoglyptus tzeltales</i>	<i>Thinodytes cyzicus</i>
<i>Chrysocharis vonones</i>	<i>Diglyphus websteri</i>	<i>Oenanogastra</i> sp.	<i>Thinodytes</i> sp.
<i>Chrysonotomyia</i> sp.	<i>Disorygma pacifica</i>	<i>Opius caricivora</i>	<i>Tribliographa</i> sp.
<i>Cirrospilus ambiguus</i>	<i>Epiclerus</i> sp.	<i>Opius chromatomyiae</i>	<i>Trichomalopsis</i> sp.
<i>Cirrospilus vittatus</i>	<i>Ganaspidium</i> sp.	<i>Opius dimidiatus</i>	<i>Trichopria</i> sp.
<i>Closterocerus cinctipennis</i>	<i>Gronotoma adachiae</i>	<i>Opius dissitus</i>	<i>Zaenecia</i> sp.
<i>Closterocerus okazakii</i>	<i>Gronotoma micromorpha</i>	<i>Opius mandibularis</i>	<i>Zagrammosoma latilineatum</i>
<i>Closterocerus pulcher</i>	<i>Gronotoma</i> sp.	<i>Opius meracus</i>	<i>Zagrammosoma multileneatum</i>
<i>Closterocerus</i> sp.	<i>Halticoptera arduine</i>	<i>Opius pallipes</i>	<i>Zagrammosoma</i> sp.
<i>Dacnusa sasakawai</i>	<i>Halticoptera circulus</i>	<i>Opius scabriventris</i>	

5.4 Decision support for eradication

A summary of key activities associated with different detection scenarios of serpentine leafminer is provided in Table 10. Differing physical detection scenarios have been chosen to highlight operational differences in urban compared with agricultural and horticultural environments.

A summary of factors to be considered for eradication or alternative action is provided in Table 11.

Table 10 Recommended responses and considerations for differing scenarios of detection of serpentine leafminer

DETECTION SCENARIO	RECOMMENDED ERADICATION/ CONTROL TREATMENTS – SEE SECTION 5.3	RECOMMENDED MOVEMENT CONTROLS – SEE SECTION 5.2	RECOMMENDED SURVEILLANCE OPTIONS – SEE SECTION 4
Commercial glasshouse/ protected cropping setting	<ul style="list-style-type: none"> ➤ Destruction of hosts within glasshouses could be considered to limit food sources or refuges (see Section 5.3.4). ➤ Chemical sprays to eradicate larvae and adults (see Section 5.3.3). ➤ Surveillance within 1 km radius of the detection and in glasshouses managed by the same business as the infested glasshouse (use visual inspections, sweep netting and yellow sticky traps for surveillance). 	<ul style="list-style-type: none"> ➤ No movement of equipment or produce without permit. ➤ Permits to include information on inspection, decontamination or destruction requirements. 	<ul style="list-style-type: none"> ➤ Minimum 1 km surveillance zone around the Infected Premises (IP). ➤ Focus on known hosts. ➤ Surveillance at linked properties. ➤ See Section 4 for points to consider when establishing a surveillance program. Key points for glasshouses are: <ul style="list-style-type: none"> • Undertake visual surveillance of all host plants. • Consider using visual surveys, sweep nets and yellow sticky traps.
Open agricultural/ horticultural setting	<ul style="list-style-type: none"> ➤ Spray infested area using appropriate insecticide (see Section 5.3.3). ➤ Surveillance within 1 km radius of the detection and survey jointly managed properties (use visual inspections, sweep netting and yellow sticky traps for surveillance). 	<ul style="list-style-type: none"> ➤ No movement of equipment or produce without permit. ➤ Permits to include information on inspection, decontamination or destruction requirements. 	<ul style="list-style-type: none"> ➤ Minimum 1 km surveillance zone around each IP. ➤ Yellow sticky traps, visual inspections and sweep netting should be undertaken in the 1 km zone. ➤ General surveillance awareness campaigns may be beneficial. ➤ Surveillance at linked properties should occur.

DETECTION SCENARIO	RECOMMENDED ERADICATION/ CONTROL TREATMENTS – SEE SECTION 5.3	RECOMMENDED MOVEMENT CONTROLS – SEE SECTION 5.2	RECOMMENDED SURVEILLANCE OPTIONS – SEE SECTION 4
Urban environment	<ul style="list-style-type: none"> ➤ Chemical sprays to eradicate larvae and adults. ➤ Surveillance within 1 km radius of the detection (use visual inspections, sweep netting and yellow sticky traps for surveillance). 	<ul style="list-style-type: none"> ➤ Movement controls in infested areas could be undertaken by permit or as voluntary movement controls (coupled with awareness campaign for urban areas). 	<ul style="list-style-type: none"> ➤ Surveillance on host plants within 1 km zone around IP (noting leaf mines, feeding and oviposition punctures are the most easily seen plant symptoms). ➤ Surveillance to be undertaken in conjunction with awareness campaigns for households providing information on the pest and symptoms. ➤ Yellow sticky traps, visual inspections and sweep netting should be undertaken in the 1 km zone. ➤ Surveillance at linked properties.
Open natural environment (e.g. detection on roadsides or national park)	<ul style="list-style-type: none"> ➤ Treatment of detections in a 1 km zone pending delimiting surveillance. ➤ Surveillance within 1 km radius of the detection (use visual inspections, sweep netting and yellow sticky traps for surveillance). 	<ul style="list-style-type: none"> ➤ Movement controls in infested areas could be undertaken by permit or as voluntary movement controls. ➤ Movement restrictions may be considered pending outcomes of delimiting surveillance. 	<ul style="list-style-type: none"> ➤ Surveillance on host plants within 1 km zone around IP (noting leaf mines, feeding and oviposition punctures are the most easily seen plant symptoms). ➤ Yellow sticky traps, visual inspections and sweep netting should be undertaken in the 1 km zone. ➤ General surveillance and awareness campaigns should be considered. ➤ Surveillance should be undertaken at linked properties.

Table 11 Summary of factors to be considered in determining whether eradication or alternative action will be taken for an incursion of serpentine leafminer

FACTORS TO CONSIDER REGARDING THE TECHNICAL FEASIBILITY OF ERADICATION
<ul style="list-style-type: none"> • The population size and population structure associated with the initial detection. • The cost effectiveness of recommended control technique options. <ul style="list-style-type: none"> - Multiple applications of several different pesticides over a period of time (weeks/months) may be required to successfully eradicate serpentine leafminer populations. • The ability to remove or destroy all serpentine leafminers by the recommended control techniques. <ul style="list-style-type: none"> - Permission to enter private premises for surveillance and treatment must be considered. - Determination of whether treatments can effectively eradicate populations within premises or environments. • The recommended control techniques are publicly acceptable. <ul style="list-style-type: none"> - Chemical treatments in residences or backyards may not be acceptable or will require negotiation with residents. • Whether emergency containment measures can effectively be put in place to contain the outbreak. • Whether there are control methods, commonly employed for endemic pests, that may prevent the establishment of, or be impacted by the treatment, of serpentine leafminer. • Legislative impediments to undertaking an eradication response. <ul style="list-style-type: none"> - If not a prescribed pest, confirmatory diagnosis will be required under legislation in some jurisdictions before powers to enter premises and undertake treatments can be enacted. - Delimiting surveillance in urban environments may be restricted to front yards and environs rather than entry into premises (unless permission from the resident can be negotiated). - Emergency permits will be required for chemical treatments. Amendments to chemical labels will be required for longer term management of the pest if eradication is not possible. • The ability to delimit the known area of infestation. <ul style="list-style-type: none"> - Determination of linked properties in an urban detection may be unfeasible given the ability of adult leafminers to disperse at a local level. High risk linkages such as the recent purchase of seedlings, cut flowers or other commodities that could vector the pest should be investigated. • The ability to identify and close the pathway for entry of the pest into Australia. <ul style="list-style-type: none"> - Pathways into Australia such as on nursery plants from overseas are managed via Post Entry Quarantine. - Natural dispersal is a possible pathway from neighbouring countries into northern Australia. Northern Australia Quarantine Strategy (NAQS) surveys offer an early detection mechanism to detect natural introductions from the north. If an outbreak is detected the pathway will need to be identified and closed. • The dispersal ability of the pest. <ul style="list-style-type: none"> - Serpentine leafminer has the ability to naturally spread over short distances through flight or wind assisted dispersal. - Nursery material that is a host of serpentine leafminer (including containerized plants, seedlings, bare rooted plants, etc.; see Appendix 1 for host list) and cut flowers that are host plants are the most likely means of spread to new areas. They may carry all life stages of the pest and will need to be managed to reduce the spread of the pest to new areas. Nursery material and cut flowers that are not a host of serpentine leafminer are at significantly lower risk (but may still transport adults across short periods if they have been in close proximity with host plants). - Plant material (potentially including fresh hay and crop debris) and soil could also facilitate the spread of the pest (via adults and pupae) to new areas. Tools, equipment, machinery and vehicles are lower risk but may still transport hitchhiking adults or pupae via soil deposits.

- Serpentine leafminer is considered unlikely to travel with harvested fruit (without leaves) as the process of harvesting or moving produce is likely to cause adults to disperse and provided soil and leaf material is absent it is unlikely to spread larvae and pupae. Leafminer are also not expected to travel via the movement of seed.
- Fresh green or leafy vegetables or fresh legume vegetables (e.g. snow peas) may allow the spread of eggs and larvae and potentially adults and pupae. Therefore, movement restrictions or mandatory treatments may be needed for some vegetable crops.
- See Table 3 for further details on possible dispersal pathways
- **The capability to detect *Liriomyza* leafminers at very low densities for the purpose of declaring freedom, and that all sites affected by the pest have or can be found.**
 - Surveillance typically relies on the use of visual inspection of host plants for mining activity as well as the use of yellow sticky traps and sweep netting.
- **The ability to put into place surveillance to confirm proof of freedom.**
 - Surveillance options are available for serpentine leafminer and include visual inspection of host plants for mines, yellow sticky traps and sweep netting.
- **Whether community consultation activities have or will be undertaken.**
 - In an urban environment, community consultation will be critical to assist in securing public support for delimiting surveillance and an eradication program.
 - Given the symptoms caused by this pest on a wide range of plants, general surveillance activities should be implemented comprising of awareness material, media releases, web-based reporting tools and the Exotic Plant Pest Hotline (1800 084 881).

6. EMERGENCY PLANT PEST RESPONSE DEED AND PLANTPLAN

The Emergency Plant Pest Response Deed (EPPRD) and PLANTPLAN should be referred to, in conjunction with this contingency plan when developing a Response Plan in the event of a pea leafminer detection in Australia.

6.1 Emergency Plant Pest Response Deed

PHA is the custodian of the EPPRD. This is a formal legally binding agreement between PHA, the Australian Government, all state and territory governments and national plant industry body signatories.

It covers the management and funding of responses to Emergency Plant Pest (EPP) incidents, including the potential for Owner Reimbursement Costs (ORCs) for growers. It also formalises the role of plant industries' participation in decision making, as well as their contribution towards the costs related to approved responses.

Under the EPPRD an EPP is defined as a pest that meets one or more of the following criteria:

- a) It is a known exotic Plant Pest the economic consequences of an occurrence of which would be economically or otherwise harmful for Australia, and for which it is considered to be in the regional and national interest to be free of the Plant Pest.
- b) It is a variant form of an established Plant Pest which can be distinguished by appropriate investigative and diagnostic methods and which, if established in Australia, would have a regional and national impact.
- c) It is a serious Plant Pest of unknown or uncertain origin which may, on the evidence available at the time, be an entirely new Plant Pest or one not listed in Schedule 13 and which if established in Australia is considered likely to have an adverse economic impact regionally and nationally.
- d) It is a Plant Pest already found in Australia that:
 - (i) is restricted to a defined area through the use of regulatory measures intended to prevent further spread of the pest out of the defined area or into an endangered area; and
 - (ii) has been detected outside the defined area; and
 - (iii) is not a native of Australia; and
 - (iv) is not the subject of any instrument for management which is agreed to be effective risk mitigation and management at a national level; and
 - (v) is considered likely to have an adverse economic impact such that an emergency response is required to prevent an incident of regional and national importance.

See: www.planthealthaustralia.com.au/biosecurity/emergency-plant-pest-response-deed/ for more information and the most recent version of the EPPRD.

6.2 PLANTPLAN

Underpinning the EPPRD is PLANTPLAN, the agreed technical response plan for an EPP incident. It provides nationally consistent guidelines for response procedures, outlining the phases of an incursion, as well as the key roles and responsibilities of industry and government during each of the phases.

See: www.planthealthaustralia.com.au/biosecurity/emergency-plant-pest-response-deed/ for more information and the most recent version of PLANTPLAN.

7. REFERENCES

- Abou-Fakhr Hammad EM, Nemer NM (2000) Population densities, spatial pattern and development of the pea leafminer (Diptera: Agromyzidae) on cucumber, swiss chard and bean. *The Journal of Agricultural Science* 134: 61–68.
- Andersen A, Nordhus E, Thanh VT, An TTT, Hung HQ, Hofsvang T (2002) Polyphagous *Liriomyza* species (Diptera: Agromyzidae) in vegetables in Vietnam. *Tropical Agriculture*, 79: 241–246.
- Andrade ME, Briceño JA, de Hoyos PM, Jiménez J (1989) Busqueda y reconocimiento de los enemigos naturales y hospedantes alternos de las principales plagas. en flores bajo invernadero en la sabana de bogota. *Acta Biológica Colombiana* 1: 45–57.
- Ávalos DS, Ricobelli G, Palacios SM, Defagó MT (2013) Evaluación de la preferencia de *Liriomyza* spp. En genotipos diferentes de garbanzo y efecto del extracto de *Melia azedarach* L.: Resultados preliminares. *Revista de la Facultad de Ciencias Agrarias*, 45:65–73.
- Bahlai CA, Goodfellow SA, Stanley-horn DE, Hallett RH (2006) Endoparasitoid Assemblage of the Pea Leafminer, *Liriomyza huidobrensis* (Diptera: Agromyzidae), in Southern Ontario. *Environmental Entomology*, 35: 351–357.
- CABI and EPPO (date unknown) Data sheets on Quarantine Pests: *Liriomyza huidobrensis*. Available from: <https://gd.eppo.int/taxon/LIRIHU/documents>.
- Calabretta C, Calabro M, Colombo A, Campo G (1995) [Spreading of *Liriomyza huidobrensis* (Blanchard)(Diptera, Agromyzidae) in protected cultures in Sicily]. [Italian]. *Informatore Fitopatologico*, 45: 24–30.
- Capinha C, Essl F, Seebens H, Moser D, Pereira HM (2015). The dispersal of alien species redefines biogeography in the Anthropocene. *Science*, 348: 1248–1251.
- Cesar Australia (2018) Prospects for biological control of the Vegetable leafminer, *Liriomyza sativae*, in Australia. Report developed as part of MT16004, Hort Innovation.
- Chavez GL and Raman KV (1987) Evaluation of trapping and trap types to reduce damage to potatoes by the leafminer, *Liriomyza huidobrensis* (Diptera, Agromyzidae). *International Journal of Tropical Insect Science*, 8: 369–372.
- Cheng L (1976) Marine Insects. North Holland Publishing Company, Amsterdam.
- Collins DW (1996) The separation of *Liriomyza huidobrensis* (Diptera: Agromyzidae) from related indigenous and non-indigenous species encountered in the United Kingdom using cellulose acetate electrophoresis. *Annals of Applied Biology*, 128: 387–398.
- Cuthbertson AGS, Mills K, Woner-Smith T, Blackburn LF, Mathers JJ and Northing P (2013) Environmental evaluation of alternative chemicals to methyl bromide for fumigation of quarantine pests in transit. *International Journal of Environmental Science and Technology*, 10: 1057–1066.
- Dawah HA and Deeming JC (2002) *Liriomyza huidobrensis* (Blanchard) (Dipt., Agromyzidae) in Arabia. *Entomologists Monthly Magazine* 138: 1656–1659.
- DAWR (2016) Leaf miner. Available from: www.agriculture.gov.au/pests-diseases-weeds/plant/leaf-miner.
- De Clercq R, Casteels H (1992) Occurrence and host plants of *Frankliniella occidentalis* and *Liriomyza huidobrensis* in Belgium 1. *EPPO Bulletin* 22: 363–366.
- De Goffau LJW (1991) *Liriomyza huidobrensis* (Blanchard) (Diptera:Agromyzidae) A new economically important leaf miner in the Netherlands. *Proceedings of the Section Experimental and Applied Entomology of the Netherlands Entomological Society* 2: 41–45.
- Dobrosavljevic J, Markovic C, Bojic S (2017) Overview of leaf miner fauna in Serbia. In: VIII International Scientific Agriculture Symposium, Jahorina, Bosnia and Herzegovina, October 2017. Faculty of Agriculture, University of East Sarajevo. Pp 1490-1498.

- EFSA Panel on Plant Health (2012) Scientific Opinion on the risks to plant health posed by *Liriomyza huidobrensis* (Blanchard) and *Liriomyza trifolii* (Burgess) to the EU territory with the identification and evaluation of risk reduction options. *EFSA Journal* 10: 3028.
- EPPO (2005) Diagnostics: *Liriomyza* spp. PM 7/53(1). *OEPP/EPPO Bulletin*, 35: 335-344.
- EPPO (2009) Disinfection of production site against *Liriomyza sativae*. *OEPP/EPPO Bulletin*, 39: 480-481.
- Fenoglio MS, Videla M, Salvo A and Morales JM (2019) Dispersal of the pea leaf miner *Liriomyza huidobrensis* (Blanchard, 1926) (Diptera: Agromyzidae): a field experiment. *Revista de la Facultad de Ciencias Agrarias UNCuyo*, 51: 343-352.
- Fisher N, Ubaidillah R, Reina P, La Salle J (2005) *Liriomyza* parasitoids of Southeast Asia. Melbourne, Australia, CSIRO. Available at www.ento.csiro.au/science/Liriomyza_ver3/index.html.
- Foba CN, Salifu D, Lagat ZO, Gitonga LM, Akutse KS, Fiaboe KK (2015) Species composition, distribution, and seasonal abundance of *Liriomyza* leafminers (Diptera: Agromyzidae) under different vegetable production systems and agroecological zones in Kenya. *Environmental Entomology* 44: 223-232.
- Foster RE (1986) Monitoring populations of *Liriomyza trifolii* (Diptera: Agromyzidae) in celery with pupal counts. *Florida Entomologist*, 69: 292-298.
- Glick PA (1960) Collecting insects by airplane with special reference to dispersal of the potato leafhopper. Technical Bulletin No. 1222. United States Department of Agriculture.
- Griffiths GCD (1962) Breeding leaf-mining flies and their parasites. *Entomologist's Record and Journal of Variation*, 74: 178-185, 203-206.
- Guantai MM, Ogol CPKO, Salifu D, Kasina JM, Akutse KS, Fiaboe KKM (2015) Differential effects of pesticide applications on *Liriomyza huidobrensis* (Diptera: Agromyzidae) and its parasitoids on pea in Central Kenya. *Journal of Economic Entomology*, 108: 662-671.
- Hanafi A (2005) Invasive pests and diseases: a challenge to IPM in greenhouse crops. *Phytoparasitica*. 33: 423-426.
- Hardy AC and Milne PS (1938) Studies in the distribution of insects by aerial currents. *The Journal of Animal Ecology*, 7: 199-229.
- He CX, Wu WW, Wang SF and Wang LZ (2001) Host plants and feeding preference of *Liriomyza huidobrensis*. *Acta Entomologica Sinica*, 44: 384-388.
- He L, Zhang Y, Xiao N, Wei J and Kuang R (2002) *Liriomyza huidobrensis* in Yunnan, China: current distribution and genetic structure of a recently established population. *Entomologia Experimentalis et Applicata*, 102: 213-219.
- He L, Zhang Y, Xiao N, Wei J, Kuang R (2002) *Liriomyza huidobrensis* in Yunnan, China: current distribution and genetic structure of a recently established population. *Entomologia Experimentalis et Applicata*, 102: 213-219.
- Heinz KM, Parrella MP, Newman JP (1992) Time-efficient use of yellow sticky traps in monitoring insect populations. *Journal of Economic Entomology*, 85: 2263-2269.
- Hidalgo JE, Carballo VM (1991) Influencia de las malezas sobre los insectos controladores naturales de *Liriomyza huidobrensis* (Blanchard), (Diptera: Agromyzidae). (The influence of weeds on the natural enemies of *Liriomyza huidobrensis* (Blanchard) (Diptera: Agromyzidae). *Manejo Integrado de Plagas* 20-21: 49-54.
- Hincapié C, Saavedra H and Tróchez P (1993) Life cycle, behaviour and natural enemies of *Liriomyza huidobrensis* (Blanchard) on bulb onion (*Allium cepa* L.). *Revista Colombiana de Entomología*, 19: 51-57.
- IPPC (2016) Diagnostic Protocol 16: Genus *Liriomyza*. Available from: www.ippc.int/static/media/files/publication/en/2017/01/DP_16_2016_En_2017-01-30.pdf.
- Iwasaki A (2004) Effect of insecticides on *Liriomyza huidobrensis* (Blanchard)(Diptera: Agromyzidae) in Hokkaido [Japan]. *Annual Report of the Society of Plant Protection of North Japan* 55: 236-239.
- Jones VP and Parrella MP (1986) The movement and dispersal of *Liriomyza trifolii* (Diptera: Agromyzidae)

in a chrysanthemum greenhouse. *Annals of applied biology*, 109: 33-39.

- Jovicich E (2009) Leafminer pest-generic incursion management plan for the Australian vegetable industry. Horticulture Australia Limited, Project: VG061113. Available from: <https://ausveg.com.au/app/data/technical-insights/docs/VG061113.pdf>.
- Koch KC, Waterhouse DF (2000) Distribución e importancia de los artrópodos asociados a la agricultura y silvicultura en Chile. *Australian Centre for International Agricultural Research, Canberra, Australia, ACIAR Monograph* 68:234.
- Korytkowski C (1982) Contribution to the Knowledge of Agromyzidae (Diptera: Acalyptratae) in Peru (Doctoral dissertation, Thesis (Magister Scientiae, in the specialty of Entomology). National Agrarian University La Molina, Graduate Academic Program).
- Kox LFF, Van Den Beld HE, Lindhout BI, De Goffau LJW (2005) Identification of economically important *Liriomyza* species by PCR-RFLP analysis. *EPPO Bulletin* 35:79–85.
- Lanzoni A, Bazzocchi GG, Burgio G, Fiacconi MR (2002) Comparative life history of *Liriomyza trifolii* and *Liriomyza huidobrensis* (Diptera: Agromyzidae) on beans: effect of temperature on development. *Environmental Entomology* 31: 797-803.
- Larraín PS, Salas CF, Jana AC, Graña SF (2013) Major artichoke pests (cynara cardunculus sub. scolymus l.) in coquimbo region, Chile. *Acta Horticulturae* 983: 301–306.
- Larrein SP (2002) Insect and mite pest incidence on sweet pepinos *Solanum muricatum* (Ait.) cultivated in the IV Region, Chile. *Agricultura-Tecnica* 62:15–26.
- Lei ZR, Wang Y, Huang DR, Cheng DF (2002) Influence of temperature on flight potential of *Liriomyza sativae* Blanchard. *Acta Entomologica Sinica*, 45: 413-415.
- Maharjan R, Oh HW, Jung C (2014) Morphological and genetic characteristics of *Liriomyza huidobrensis* (Blanchard)(Diptera: Agromyzidae) infesting potato crops in Korea. *Journal of Asia-Pacific Entomology* 17: 281-286.
- Maharjan R, Oh HW, Jung C (2014) Morphological and genetic characteristics of *Liriomyza huidobrensis* (Blanchard) (Diptera: Agromyzidae) infesting potato crops in Korea. *Journal of Asia-Pacific Entomology*, 17: 281–286.
- Martin AD, Stanley-Horn D, Hallett RH (2006) Adult host preference and larval performance of *Liriomyza huidobrensis* (Diptera: Agromyzidae) on selected hosts. *Environmental Entomology*, 34: 1170–1177.
- Martin AD, Vernon RS, Hallett RH (2005) Influence of colour and trap height on captures of adult pea leafminer, *Liriomyza huidobrensis* (Blanchard) (Diptera: Agromyzidae), in celery. *Journal of the Entomological Society of Ontario*, 136: .25-35.
- Masetti A, Luchetti A, Mantovani B, Burgio G (2006) Polymerase chain reaction-restriction fragment length polymorphism assays to distinguish *Liriomyza huidobrensis* (Diptera: Agromyzidae) from associated species on Lettuce Cropping Systems in Italy. *Journal of Economic Entomology* 99: 1268–1272.
- Milla, K. and Reitz, S., 2005. Spatial/temporal model for survivability of pea leafminer (*Liriomyza huidobrensis*) in warm climates: a case study in south Florida, USA. *European Journal of scientific research*, 7: 65-73.
- Mujica N (2007) Malezas hospederas de moscas minadoras (Diptera: Agromyzidae) y sus parasitoides en el ecosistema de papa en La Molina. Doctoral Thesis, Lima, Perú. UNALM.
- Mujica N, Kroschel J (2011) Leafminer fly (Diptera: Agromyzidae) occurrence, distribution, and parasitoid associations in field and vegetable crops along the Peruvian coast. *Environmental Entomology* 40: 217–230.
- Mujica N, Sporleder M, Carhuapoma P, Kroschel J (2017) A temperature-dependent phenology model for *Liriomyza huidobrensis* (Diptera: Agromyzidae). *Journal of Economic Entomology*, 110: 1333-1344.
- Murphy ST and LaSalle J (1999) Balancing biological control strategies in the IPM of New World invasive *Liriomyza* leafminers in field vegetable crops. *Biocontrol News and Information*, 20: 91-104.

- Musundire R, Chabi-Olaye A, Krüger K (2012) Host plant effects on morphometric characteristics of *Liriomyza huidobrensis*, *L. sativae* and *L. trifolii* (Diptera: Agromyzidae). *Journal of Applied Entomology*, 136: 97–108.
- Nakamura S, Masuda T, Mochizuki A, Konishi K, Tokumaru S, Ueno K, Yamaguchi T (2013) (2013) Primer design for identifying economically important *Liriomyza* species (Diptera: Agromyzidae) by multiplex PCR. *Molecular Ecology Resources* 13: 96–102.
- Niño L, Prieto L, Santiago V, Acevedo E (2009) Fluctuación poblacional de la mosca minadora (*Liriomyza huidobrensis* Blanchard) en cultivos de papa de Pueblo Llano, Estado Mérida, Venezuela. *Entomotropica* 24:65–70.
- Noujeim E, Sakr J, Nemer N (2013) Potential of entomopathogenic nematodes application against *Liriomyza huidobrensis* Blanchard in Lebanon. In: Proceedings, 4th International Scientific Symposium Agrosys Bosnia and Herzegovina, Faculty of Agriculture, University of East Sarajevo. Bosnia and Herzegovina. pp 692–698.
- Okoth CA, Deng AL, Tabu IM, Akutse KS, Fiaboe KKM (2014) Effect of host plant on feeding, biological and morphological parameters of *Liriomyza huidobrensis* Blanchard (Diptera: Agromyzidae). *African Entomology*, 22: 577–588.
- Ozawa A, Saito T and Ota M (1999) Dispersal of adults of American Serpentine Leafminer, *Liriomyza trifolii* (Burgess) (Diptera: Agromyzidae) in fields. *Japanese Journal of Applied Entomology and Zoology*, 43: 49–54.
- Pang B, Gao J, Zhou X, Wang J (2006) Relationship between host plant preference of *Liriomyza huidobrensis* (Blanchard)(Diptera: Agromyzidae) and secondary plant compounds and trichomes of host foliage. *Acta Entomologica Sinica* 49: 810–815.
- PHA (2008) Threat Specific Contingency Plan: American serpentine leafminer, *Liriomyza trifolii*, bundled with *L. cicerina*, *L. huidobrensis*, *L. sativae*, *L. bryoniae* and *Chromatomyia horticola*. Available from: www.planthealthaustralia.com.au/wp-content/uploads/2013/03/American-serpentine-and-other-leaf-miners-CP-2008.pdf.
- PHA (2015) PLANTPLAN Guideline document: Disinfection and Decontamination. available from: www.planthealthaustralia.com.au/wp-content/uploads/2015/12/Guidelines-Disinfection-and-decontamination.pdf.
- Pirtle EI, Van Rooyen A, Maino J, Weeks AR, Umina PA (*in press*) A molecular model for biomonitoring of an exotic plant-pest: leafmining for environmental DNA.
- Queensland Herbarium, (2016). Collection and preserving plant specimens, a manual. 2nd edition. Department of Science, Information Technology and Innovation, Brisbane. Available from: www.qld.gov.au/data/assets/pdf_file/0032/67469/collecting-manual.pdf.
- Rauf A, Merle Shepard B, Johnson MW (2000) Leafminers in vegetables, ornamental plants and weeds in Indonesia: Surveys of host crops, species composition and parasitoids. *International Journal of Pest Management* 46: 257–266.
- Saethre M (1996) Pest risk assessment (PRA) for the South American leafminer *Liriomyza huidobrensis*. Planteforsk, Norwegian Agricultural Inspection Service. Available from: <https://core.ac.uk/reader/285995081>.
- Salvo A, Valladares G (2002) Plant-Related Intraspecific Size Variation in Parasitoids (Hymenoptera: Parasitica) of a Polyphagous Leafminer (Diptera: Agromyzidae). *Environmental Entomology* 31:874–879.
- Salvo A, Valladares GR (1997) An analysis of leaf-miner and plant host ranges of three Chrysocharis species (Hym.: Eulophidae) from Argentina. *Entomophaga* 42:417–426.
- Scheffer SJ and Lewis ML (2001) Two nuclear genes confirm mitochondrial evidence of cryptic species within *Liriomyza huidobrensis* (Diptera: Agromyzidae). *Annals of the Entomological Society of America*, 94: 648–653.
- Scheffer SJ, Lewis ML, Joshi RC (2006) DNA Barcoding Applied to Invasive Leafminers (Diptera: Agromyzidae) in the Philippines. *Annals of the Entomological Society of America*, 99: 204–210.

- Scheffer SJ, Wijesekara A, Visser D, Hallett RH (2001) Polymerase chain reaction-restriction fragment-length polymorphism method to distinguish *Liriomyza huidobrensis* from *L. langei* (Diptera: Agromyzidae) applied to three recent leafminer invasions. *Journal of Economic Entomology*, 94: 1177-1182.
- Shepard BM and Braun SAR (1998) Seasonal incidence of *Liriomyza huidobrensis* (Diptera: Agromyzidae) and its parasitoids on vegetables in Indonesia. *International Journal of Pest Management*, 44: 43-47.
- Shepard BM, Samsudin, Braun AR (1998) Seasonal incidence of *Liriomyza huidobrensis* (Diptera: Agromyzidae) and its parasitoids on vegetables in Indonesia. *International Journal of Pest Management* 44: 43-47.
- Shiao SF and Wu WJ (2000) *Liriomyza huidobrensis* (Blanchard), a newly invaded insect of economic importance to Taiwan (Diptera: Agromyzidae). *Plant Protection Bulletin Taipei*, 42: 249-254.
- Shindo J (2006) Occurrence of pea Leafminer, *Liriomyza huidobrensis* (Blanchard) (Diptera: Agromyzidae), in Aomori Prefecture. *Annual Report of the Society of Plant Protection of North Japan* 56: 188-191.
- Singh OL and Singh CJ (2013) Population Dynamics and Seasonal Incidence of *Liriomyza huidobrensis* (Blanchard) (Diptera: Agromyzidae) on Onion vegetable in Manipur. *International Journal of Advancements in Research & Technology* 2: 89-95.
- Sivapragasam A, Syed AR, La Salle J, Ruwaida M (1999) Parasitoids of invasive agromyzid leafminers on vegetables in Peninsular Malaysia. In: Proceedings of Symposium on Biological Control in the Tropics. pp 8-19.
- Spencer KA (1973) Agromyzidae (Diptera) of Economic Importance. *Series Entomologica* 9. The Hague: W. Junk. 418 pp.
- Spencer KA (1983) Leaf mining Agromyzidae (Diptera) in Costa Rica. *Revista de Biología tropical*, 31: 41-67.
- Spencer KA (1989). Leaf miners. In Plant Protection and Quarantine, Vol. 2, Selected Pests and Pathogens of Quarantine Significance (ed Kahn RP). CRC Press, Boca Raton, pp. 77-98.
- Spencer KA (1990). Host specialization in the world Agromyzidae (Diptera). *Series Entomologica* 45. Kluwer Academic Publishers, Dordrecht. 444 pp.
- Spencer KA, Martínez, M, Étienne J (1992) The Agromyzidae (Diptera) of Guadeloupe. *Annales de la Société Entomologique de France* 28: 251-302.
- Steck G (1999) Pest alert: pea leaf miner, *Liriomyza huidobrensis* (Diptera: Agromyzidae). Florida Department of Agriculture and Consumer Services, Division of Plant Industry. Publication number: DACS-P-01691. Available from: www.freshfromflorida.com/content/download/66426/1601319/Pest_Alert_-_Liriomyza_huidobrensis_Pea_leafminer.pdf.
- Suss, L. (1991) First record in Italy of *Liriomyza huidobrensis*. *Bollettino di Zoologia Agraria e di Bachicoltura* 23, 197-202.
- Takano SI, Iwaizumi R, Nakanishi Y, Someya H (2008) Laboratory hybridization between the two clades of *Liriomyza huidobrensis* (Diptera: Agromyzidae). *Applied Entomology and Zoology*, 43: 397-402.
- TCR White (1970) Airborne arthropods collected in south Australia with a drogue-net towed by a light aircraft *Pacific Insects* 12: 251-259.
- Tran DH and Takagi M (2005) Susceptibility of the stone leek leafminer *Liriomyza chinensis* (Diptera: Agromyzidae) to insecticides. *Journal of the Faculty of Agriculture, Kyushu University*, 50: 383-390.
- Tran DH (2009) Agromyzid leafminers and their parasitoids on vegetables in central Vietnam. *Journal of the International Society for Southeast Asian Agricultural Sciences* 15: 21-33.
- Trouvé C, Martinez M, Phalip M, Martin C (1991) A new pest in Europe, the South American miner fly. *Phytoma* 429: 42-46.
- Trumble JT and Nakakihara H (1983) Occurrence, parasitization, and sampling of *Liriomyza* species (Diptera: Agromyzidae) infesting celery in California. *Environmental Entomology*, 12: 810-814.

- Valladares (1984) Sobre el Género *Liriomyza* Mik 1894 (Diptera, Agromyzidae) en la República Argentina. *Revista de la Sociedad Entomológica Argentina* 43: 13–36.
- Valladares G, Pinta D, Salvo A (1996) La mosca minadora *Liriomyza huidobrensis* Diptera, Agromyzidae en cultivos de Córdoba. *Hortic Argentina* 15: 13–18.
- Valladares G, Salvo A, Saini E (2011) Moscas minadoras del girasol y sus enemigos naturales. *Revista de Investigaciones Agropecuarias* 37: 180–188.
- Valladares GR, Salvo A, Videla M (1999) Moscas minadoras en cultivos de Argentina. *Horticulture Argentina* 18: 44–45.
- Venette RC, Hutchison WD, Burkness EC, O'Rourke PK (1980) Alfalfa blotch leafminer: Research update. University of Minnesota. Available from: <https://ipmworld.umn.edu/venette>.
- Vercambre B, A De Crozals (1993) Relationship between *Liriomyza huidobrensis* and *L. trifolii* in Reunion Island. In: *Liriomyza: Conference on leaf-mining flies in cultivated plants*, 24–26 March 1993, Montpellier (France), pp. 71–78.
- Wei J, Zou L, Kuang R, He L (2000) Influence of leaf tissue structure on host feeding selection by pea leafminer *Liriomyza huidobrensis* (Diptera: Agromyzidae). *Zoological Studies-Taipe* 39:295–300.
- Weintraub PG (2001) Changes in the dynamics of the leafminer, *Liriomyza huidobrensis*, in Israeli potato fields. *International Journal of Pest Management*, 47: 95–102.
- Weintraub PG and Horowitz AR (1995) The newest leafminer pest in Israel, *Liriomyza huidobrensis*. *Phytoparasitica*, 23:177–184.
- Weintraub PG and Horowitz AR (1996) Spatial and diel activity of the pea leafminer (Diptera: Agromyzidae) in potatoes, *Solanum tuberosum*. *Environmental Entomology*, 25: 722–726.
- Weintraub PG, Scheffer SJ, Visser D, et al (2017) The invasive *Liriomyza huidobrensis* (Diptera: Agromyzidae): understanding its pest status and management globally. *Journal of Insect Science*, 17: issue 1.
- Weintraub PG, Scheffer SJ, Visser D, Valladares G, Correa AS, Shepard BM, Rauf A, Murphy ST, Mujica N, MacVean C, Kroschel J, Kishinevsky M, Joshi RC, Johansen NS, Hallett RH, Civelek HS, Chen B, Metzler HB (2017) The invasive *Liriomyza huidobrensis* (Diptera: Agromyzidae): understanding its pest status and management globally. *Journal of Insect Science*, 28: 1–27.
- Wichmann MC, Alexander MJ, Soons MB, Galsworthy S, Dunne L, Gould R, Fairfax C, Niggemann M, Hails RS and Bullock JM (2009) Human-mediated dispersal of seeds over long distances. *Proceedings of the Royal Society B: Biological Sciences*, 276: 523–532.
- Yábar E, Gianoli E, Echegaray ER (2002) Insect pests and natural enemies in two varieties of quinoa (*Chenopodium quinoa*) at Cusco, Peru. *Journal of Applied Entomology*, 126: 275–280.
- Yoshimoto CM and Gressitt JL (1964). Dispersal studies on Aphididae, Agromyzidae and Cynipoidea. *Pacific Insects*, 6: 525–531.
- Zehnder GW and Trumble JT (1984) Intercrop movement of leafminers. *California Agriculture*, 38: 7–8.
- Zoebisch TG and Schuster DJ (1990) Influence of height of yellow sticky cards on captures of adult leafminer (*Liriomyza trifolii*) (Diptera: Agromyzidae) in staked tomatoes. *The Florida Entomologist*, 73: 505–507.
- Addendum:
- Gitonga ZM, Chabi-Olaye A, Mithöfer D, Okello JJ & Ritho CN. 2010. Control of invasive *Liriomyza* leafminer species and compliance with food safety standards by small scale snow pea farmers in Kenya. *Crop Protection* 298: 1472–1477.

8.APPENDICES

Appendix 1: Known hosts of serpentine leafminer

The serpentine leafminer (*Liriomyza huidobrensis*) is highly polyphagous, with a wide host range across at least 15 families (Table 12). While host records for *L. huidobrensis* have been reviewed in previous studies (Weintraub et al. 2017), in order to facilitate trade and market access decisions, a thorough review was conducted as part of the Horticulture Innovation Australia funded Project MT16004 (R&E program for control, eradication and preparedness for vegetable leafminer), with a focus on evidence for lifecycle completion in the field. The results of this review, including commercial, ornamental and non-cultivated hosts of serpentine leafminer (*L. huidobrensis*), are presented in Table 13, Table 14 and Table 15 below. No guarantee can be made that these lists are exhaustive. Each record was scrutinized based on evidence provided showing completion of lifecycle within the field, and then included within the appropriate table. It is important to note that non-cultivated plants often serve as important reservoirs for *Liriomyza* pests (e.g. Schuster et al. 1991), and upon incursion into a new region, host ranges may expand to include native plant species. Moreover, host preferences may vary considerably across populations. For example, although chili (*Capsicum* spp.) is widely reported internationally as a favoured host of the closely related agromyzid *Liriomyza sativae*, the population of *L. sativae* which has established within the Torres Strait (IPPC 2017) has very rarely been observed to attack chili (despite it being an abundant garden host in the region) (Elia Pirtle pers. comm).

Number of host records for *L. huidobrensis* within plant families

Only records with either comprehensive or partial evidence of lifecycle completion in the field are included (see Table 13 and Table 14). Multiple records of the same species are included. Thus, the number of records does not indicate total number of affected species within each family. *Liriomyza huidobrensis* is highly polyphagous and no guarantee can be made that this list is exhaustive.

Table 12 Number of host records for *L. huidobrensis* within plant families.

Family	Number of records
Asteraceae	81
Brassicaceae	36
Fabaceae	32
Solanaceae	29
Amaranthaceae	27
Cucurbitaceae	20
Caryophyllaceae	13
Apiaceae	12
Amaryllidaceae	10
Malvaceae	7
Ranunculaceae	7
Poaceae	6
Convolvulaceae	5
Lamiaceae	5

Family	Number of records
Polygonaceae	5
Chenopodiaceae	4
Gentianaceae	4
Passifloraceae	1
Plantaginaceae	4
Plumbaginaceae	3
Tropaeolaceae	3
Violaceae	3
Alstroemeriaceae	2
Araceae	2
Araliaceae	2
Balsaminaceae	2
Basellaceae	2
Campanulaceae	2
Euphorbiaceae	2
Iridaceae	2
Liliaceae	2
Primulaceae	2
Verbenaceae	2
Acanthaceae	1
Alismataceae	1
Apocynaceae	1
Asparagaceae	1
Asphodelaceae	1
Calceolariaceae	1
Hydrangeaceae	1
Linaceae	1
Linderniaceae	1
Menispermaceae	1
Onagraceae	1
Onagroideae	1
Oxalidaceae	1
Papaveraceae	1
Polemoniaceae	1
Rosaceae	1
Scrophulariaceae	1

Commercial, ornamental and non-cultivated plants for which comprehensive evidence of lifecycle completion in the field for the serpentine leafminer (*Liriomyza huidobrensis*) has been provided within the scientific literature

This includes all records where the following conditions were met:

- (1) Adults were reared from plant material infested with eggs or from larvae collected in the field; and
- (2) Emerging adults were subsequently identified via morphology or molecular diagnostics; and
- (3) Scientific names of hosts are reported

In addition, all records from Kenneth Spencer, the foremost authority on leafminer taxonomy, are included. Records for plants identified only to the genus level (sp. or spp.) that did not meet the above conditions are included only if there are other records for species within this genus that met both conditions. Records for specific varieties within a species that did not meet the above conditions are included only if there are other records confirming lifecycle completion for that same species that met both conditions. Scientific names in the table appear as they were originally cited. A number of these names are now recognised as synonyms, however for brevity, we are not reporting the currently accepted names for these taxa (refer to WorldFloraOnline.org, Australian Plant Census, or Catalogue of Life for current taxonomic decisions). All common names associated with each scientific name, across all reports, are included. For all non-cultivated plants, and asterisk after the scientific name indicates that the Atlas of Living Australia reports this species present within Australia (Atlas of Living Australia 2020). Additionally, many plants labelled as commercial and ornamental have escaped cultivation and naturalised in Australia. Records that report only partial confirmations of lifecycle completion are included in Table 14, while unverified records are included in Table 15. *Liriomyza huidobrensis* is highly polyphagous and no guarantee can be made that this list is exhaustive.

Table 13 Commercial, ornamental and non-cultivated plants for which comprehensive evidence of lifecycle completion in the field for the serpentine leafminer (*Liriomyza huidobrensis*) has been provided within the scientific literature.

Host type	Common names	Scientific name	Variety	Family	Records with comprehensive evidence	All other records (partial and unverified records)
Commercial		<i>Beta</i> sp.		Amaranthaceae		Collins (1996)
Commercial	Beet; Chard; Beetroot	<i>Beta vulgaris</i>		Amaranthaceae	Nakamura et al. (2013); Rauf et al. (2000); Scheffer et al. (2001); Takano et al. (2008)	Andersen et al. (2008); Echevarria et al. (1994); He et al. (2002); (Koch and Waterhouse 2000); Korytkowski (2014); Korytkowski (1982); Ripa et al. (1995); Saunders et al. (1998); Spencer (1973); Weintraub et al. (2017); Wijesekara (2010); Kox et al. (2005)

Host type	Common names	Scientific name	Variety	Family	Records with comprehensive evidence	All other records (partial and unverified records)
Commercial	Swisschard; Chard; Common beet	<i>Beta vulgaris</i>	<i>cicla</i>	Amaranthaceae	Hammad and Nemer (2000); Mujica and Kroschel (2011); Salvo and Valladares (2002)	He et al. (2001); He et al. (2002); Salvo and Valladares (1997); Valladares (1984); Valladares et al. (1996); Valladares et al. (1999); Valladares et al. (2011)
Commercial	Beetroot	<i>Beta vulgaris</i>	<i>rapacea</i>	Amaranthaceae	Salvo and Valladares (2002)	Valladares (1984); Valladares et al. (1996); Valladares et al. (1999); Valladares et al. (2011)
Commercial		<i>Beta vulgaris</i>	<i>rubra</i>	Amaranthaceae		Shiao and Wu (2000)
Commercial	Beetroot	<i>Beta vulgaris</i>	<i>saccharifera</i>	Amaranthaceae		Hidalgo and Carballo (1991)
Commercial	Beetroot	<i>Beta vulgaris</i>	<i>vulgaris</i>	Amaranthaceae	Mujica and Kroschel (2011)	
Commercial	Spinach; Silverbeet	<i>Spinacia oleracea</i>		Amaranthaceae	Foba et al. (2015); Mujica and Kroschel (2011); Nakamura et al. (2013); Rauf et al. (2000)	Andersen et al. (2008); Bahlai et al. (2006); He et al. (2001); (Koch and Waterhouse 2000); Korytkowski (1982); Martin et al. (2005); Spencer (1973); Valladares et al. (1996); Kox et al. (2005)
Commercial	Leek; Leeks	<i>Allium ampeloprasum</i>		Amaryllidaceae	Rauf et al. (2000)	Hincapie et al. (1993)
Commercial	Onion; Shallot; Multiplier onion	<i>Allium cepa</i>		Amaryllidaceae	Foba et al. (2015); Rauf et al. (2000)	Andersen et al. (2002); Andersen et al. (2008); He et al. (2001); Hidalgo and Carballo (1991); Hincapie et al. (1993); Korytkowski (2014); Korytkowski (1982); Kuhnke et al. (1998); Martin et al. (2005); Pang et al. (2006); Rauf et al. (2000); Spencer (1973); Ueno (2006); Verjel-Manzano and Mejia-Florez (2000); Weintraub et al. (2017); (Koch and Waterhouse 2000)
Commercial	Scallion	<i>Allium cepa</i>	<i>aggregatum</i>	Amaryllidaceae	Mujica and Kroschel (2011)	Korytkowski (1982)
Commercial		<i>Allium cepa</i>	<i>cepa</i>	Amaryllidaceae		Scheffer et al. (2006)

Host type	Common names	Scientific name	Variety	Family	Records with comprehensive evidence	All other records (partial and unverified records)
Commercial	Japanese bunching onion; Scallion; Welsh onion	<i>Allium fistulosum</i>		Amaryllidaceae	Rauf et al. (2000)	He et al. (2001); Hidalgo and Carballo (1991); Pang et al. (2006); Shepard et al. (1998); Shiao and Wu (2000); Ueno (2006); Wei et al. (2000)
Commercial	Garlic	<i>Allium sativum</i>		Amaryllidaceae	Rauf et al. (2000)	He et al. (2001); Hincapie et al. (1993); Pang et al. (2006); Saunders et al. (1998); Silva (1993); Wei et al. (2000)
Commercial		<i>Allium</i> sp.		Amaryllidaceae	Scheffer et al. (2001)	
Commercial	Celery	<i>Apium graveolens</i>		Apiaceae	Larrain and Munoz (1997); Rauf et al. (2000); Sivapragasam and Syed (1999)	Bahlai et al. (2006); de Clercq and Casteels (1992); He et al. (2001); He et al. (2002); Hidalgo and Carballo (1991); (Koch and Waterhouse 2000); Korytkowski (2014); Korytkowski (1982); Kuhnke et al. (1998); Martin et al. (2005); Pang et al. (2006); Romero-Zuniga et al. (1991); Shepard et al. (1998); Spencer (1973); Valladares (1984); Valladares et al. (1996); Valladares et al. (1999); Valladares et al. (2011); Wei et al. (2000); Weintraub (1999); Weintraub et al. (2017)
Commercial	Celery	<i>Apium graveolens</i>	<i>dulce</i>	Apiaceae	Mujica and Kroschel (2011)	He et al. (2001); Korytkowski (2014); Scheffer et al. (2006)
Commercial		<i>Apium</i> sp.		Apiaceae		Collins (1996)
Commercial	Carrot	<i>Daucus carota</i>		Apiaceae	Rauf et al. (2000)	Korytkowski (2014); Korytkowski (1982); Shepard et al. (1998)
Commercial		<i>Daucus carota</i>	<i>sativa</i>	Apiaceae		Korytkowski (2014)
Commercial	Carrot	<i>Daucus carota</i>	<i>sativus</i>	Apiaceae		Scheffer et al. (2006)
Commercial	Lettuce; Garden	<i>Lactuca sativa</i>		Asteraceae	Masetti et al. (2006); Mujica and Kroschel	Andersen et al. (2002); Andersen et al. (2008); Bahlai et al. (2006); de Clercq

Host type	Common names	Scientific name	Variety	Family	Records with comprehensive evidence	All other records (partial and unverified records)
	lettuce				(2011); Rauf et al. (2000); Takano et al. (2008)	and Casteels (1992); Echevarria et al. (1994); Godinho and Mexia (2000); He et al. (2002); Hincapie et al. (1993); (Koch and Waterhouse 2000); Korytkowski (2014); Korytkowski (1982); Kuhnke et al. (1998); Martin et al. (2005); Olivera and Bordat (1996); Weintraub et al. (2017); Parish et al. (2017); Salvo and Valladares (1997); Spencer (1973); Takano et al. (2008); Valladares (1984); Valladares et al. (1996); Valladares et al. (1999); Valladares et al. (2011); Weintraub et al. (2017)
Commercial		<i>Lactuca sativa</i>	<i>asparagina</i>	Asteraceae		Wei et al. (2000)
Commercial		<i>Lactuca sativa</i>	<i>capitata</i>	Asteraceae		Scheffer et al. (2006); Wei et al. (2000)
Commercial		<i>Lactuca sativa</i>	<i>intybeca</i>	Asteraceae		Shiao and Wu (2000)
Commercial	Indian nightshade	<i>Basella alba</i>		Basellaceae	Rauf et al. (2000)	He et al. (2001)
Commercial	Chinese cabbage; Petsai; Common yellow mustard	<i>Brassica campestris</i>		Brassicaceae	Rauf et al. (2000); Sivapragasam et al. (1999); Sivapragasam and Syed (1999)	Hidalgo and Carballo (1991); Pang et al. (2006); Romero-Zuniga et al. (1991)
Commercial		<i>Brassica campestris</i>	<i>pekinensis</i>	Brassicaceae	Mujica and Kroschel (2011)	He et al. (2001)
Commercial	Turnip	<i>Brassica campestris</i>	<i>rapa</i>	Brassicaceae	Mujica and Kroschel (2011)	Hincapie et al. (1993)
Commercial	Pak-choi cabbage; Petsai	<i>Brassica chinensis</i>		Brassicaceae	Rauf et al. (2000); Sivapragasam et al. (1999)	Andersen et al. (2008); Shepard et al. (1998)
Commercial	Leaf mustard; Mustard greens;	<i>Brassica juncea</i>		Brassicaceae	Rauf et al. (2000); Scheffer et al. (2001)	Andersen et al. (2008); Bahlai et al. (2006); He et al. (2001); Scheffer et al.

Host type	Common names	Scientific name	Variety	Family	Records with comprehensive evidence	All other records (partial and unverified records)
	Radish; Chinese mustard; Indian mustard; Mustard					(2006); Silva (1993); Wei et al. (2000)
Commercial	Gai lan; Cabbage; Broccoli; Caisin; Cabbage; Cauliflower; Field cabbage	<i>Brassica oleracea</i>		Brassicaceae	Rauf et al. (2000); Scheffer et al. (2001)	Bahlai et al. (2006); Echevarria et al. (1994); Korytkowski (2014); Korytkowski (1982); Pang et al. (2006); Saunders et al. (1998); Shepard et al. (1998)
Commercial	Kale	<i>Brassica oleracea</i>	<i>acephala</i>	Brassicaceae	Foba et al. (2015)	He et al. (2001)
Commercial	Cauliflower	<i>Brassica oleracea</i>	<i>botrytis</i>	Brassicaceae		He et al. (2001); Korytkowski (2014)
Commercial	Cabbage	<i>Brassica oleracea</i>	<i>capitata</i>	Brassicaceae	Mujica and Kroschel (2011)	He et al. (2001); Hincapie et al. (1993); Scheffer et al. (2006); Weintraub et al. (2017)
Commercial		<i>Brassica oleracea</i>	<i>caulorapa</i>	Brassicaceae		He et al. (2001)
Commercial		<i>Brassica oleracea</i>	<i>geminifera</i>	Brassicaceae		Weintraub et al. (2017)
Commercial	Broccoli	<i>Brassica oleracea</i>	<i>italica</i>	Brassicaceae		He et al. (2001); Scheffer et al. (2006); Shepard et al. (1998); Weintraub et al. (2017)
Commercial	Turnip; Rinsho; Chinese cabbage; Bokchoy	<i>Brassica rapa</i>		Brassicaceae	Rauf et al. (2000); Takano et al. (2008)	Collins (1996); Scheffer et al. (2006); Shepard et al. (1998); Valladares et al. (1999); Valladares et al. (2011)
Commercial	Pak choi; Pechay; Chinese	<i>Brassica rapa</i>	<i>chinensis</i>	Brassicaceae	Takano et al. (2008)	Andersen et al. (2008)

Host type	Common names	Scientific name	Variety	Family	Records with comprehensive evidence	All other records (partial and unverified records)
	cabbage					
Commercial		<i>Brassica rapa</i>	<i>cultivar Ciaxin</i>	Brassicaceae		CABI (2019)
Commercial		<i>Brassica</i> sp.		Brassicaceae		Hincapie et al. (1993)
Commercial	Watercress	<i>Nasturtium officinale</i>		Brassicaceae	Rauf et al. (2000)	He et al. (2001); Weintraub et al. (2017)
Commercial		<i>Nasturtium</i> spp.		Brassicaceae		Valladares (1984)
Commercial	Radish; Chinese radish (Daikon); Wild radish; Garden radish; White radish	<i>Raphanus sativus</i>		Brassicaceae	Mujica and Kroschel (2011); Rauf et al. (2000)	He et al. (2001); Collins (1996); Echevarria et al. (1994); Shepard et al. (1998); Weintraub et al. (2017)
Commercial		<i>Raphanus sativus</i>	<i>sativus</i>	Brassicaceae		Scheffer et al. (2006)
Commercial	Sweet potato	<i>Ipomoea batatas</i>		Convolvulaceae	Rauf et al. (2000)	Shepard et al. (1998)
Commercial	Watermelon	<i>Citrullus lanatus</i>		Cucurbitaceae	Foba et al. (2015)	(Koch and Waterhouse 2000); Ripa et al. (1995)
Commercial	Cucumber	<i>Cucumis sativus</i>		Cucurbitaceae	Hammad and Nemer (2000); Masetti et al. (2006); Mujica and Kroschel (2011); Nakamura et al. (2013); Rauf et al. (2000)	Andersen et al. (2002); Weintraub et al. (2017); Hincapie et al. (1993); Kuhnke et al. (1998); Martin et al. (2005); Weintraub et al. (2017); Parish et al. (2017); Ripa et al. (1995); Shiao and Wu (2000); Valladares et al. (1996); Valladares et al. (1999); Valladares et al. (2011); Yabas et al. (1995)

Host type	Common names	Scientific name	Variety	Family	Records with comprehensive evidence	All other records (partial and unverified records)
Commercial		<i>Cucumis</i> spp.		Cucurbitaceae		(Koch and Waterhouse 2000)
Commercial	Pumpkin; Winter squash; Squash	<i>Cucurbita maxima</i>		Cucurbitaceae	Foba et al. (2015); Mujica and Kroschel (2011)	Korytkowski (2014); Korytkowski (1982); Valladares et al. (1996); Valladares et al. (1999); Valladares et al. (2011)
Commercial	Courgette	<i>Cucurbita maxima</i>	<i>zapallito</i>	Cucurbitaceae	Salvo and Valladares (2002)	Salvo and Valladares (1997); Valladares et al. (1996); Valladares et al. (1999); Valladares et al. (2011)
Commercial	Butternut squash; Pumpkin; Crookneck squash	<i>Cucurbita moschata</i>		Cucurbitaceae	Foba et al. (2015)	Pang et al. (2006); Valladares et al. (1996); Valladares et al. (1999); Valladares et al. (2011)
Commercial		<i>Cucurbita moschata</i>	<i>medullosa</i>	Cucurbitaceae		Pang et al. (2006)
Commercial	Courgette; Zucchini; Pumpkin	<i>Cucurbita pepo</i>		Cucurbitaceae	Foba et al. (2015); Mujica and Kroschel (2011); Scheffer et al. (2001)	He et al. (2001); Korytkowski (1982); Olivera and Bordat (1996); Pang et al. (2006); Spencer (1973)
Commercial		<i>Cucurbita pepo</i>	<i>pepo</i>	Cucurbitaceae		Scheffer et al. (2006)
Commercial	Squash	<i>Cucurbita</i> sp.		Cucurbitaceae		Korytkowski (2014); Salvo and Valladares (1997); Valladares (1984); Valladares et al. (1996)
Commercial	Bitter gourd; Balsam pear	<i>Momordica charantia</i>		Cucurbitaceae	Foba et al. (2015)	Pang et al. (2006)
Commercial	Swartz Chayote	<i>Sechium edule</i>		Cucurbitaceae	Rauf et al. (2000)	He et al. (2001)
Commercial	Chickpea	<i>Cicer arietinum</i>		Fabaceae	Avalos et al. (2013)	
Commercial	Soybean; Soya bean	<i>Glycine max</i>		Fabaceae	Rauf et al. (2000)	He et al. (2001); Parish et al. (2017); Valladares (1984); Valladares et al. (1999)

Host type	Common names	Scientific name	Variety	Family	Records with comprehensive evidence	All other records (partial and unverified records)
Commercial	Sweet Dolichos; Field bean	<i>Lablab purpureus</i>		Fabaceae	Foba et al. (2015)	
Commercial	Alfalfa; Lucerne	<i>Medicago sativa</i>		Fabaceae	Mujica and Kroschel (2011)	Echevarria et al. (1994); Gloria and Salas (1976); (Koch and Waterhouse 2000); Korytkowski (2014); Valladares (1984); Valladares et al. (1999); Valladares et al. (2011)
Commercial	Sieva; Lima bean; Lima bean	<i>Phaseolus lunatus</i>		Fabaceae	Sivapragasam et al. (1999)	
Commercial	Beans; Grean beans; Pole beans	<i>Phaseolus</i> sp.		Fabaceae		Scheffer et al. (2006)
Commercial	Green bean; French bean; Kidney bean; Bean; Common bean; Snap bean	<i>Phaseolus vulgaris</i>		Fabaceae	Foba et al. (2015); Larrain and Munoz (1997); Mujica and Kroschel (2011); Rauf et al. (2000); de Souza (1986); Hammad and Nemer (2000); Salvo and Valladares (2002)	Andersen et al. (2002); Andersen et al. (2008); Masetti et al. (2006); Olivera and Bordat (1996); Pang et al. (2006); Parish et al. (2017); Saunders et al. (1998); Verjel-Manzano and Mejia-Florez (2000); Echevarria et al. (1994); Godinho and Mexia (2000); Weintraub et al. (2017); He et al. (2001); He et al. (2002); Hidalgo and Carballo (1991); Hincapie et al. (1993); Iwasaki (2004b); Korytkowski (2014); Korytkowski (1982);

Host type	Common names	Scientific name	Variety	Family	Records with comprehensive evidence	All other records (partial and unverified records)
						Macdonald et al. (2003); Musundire et al. (2012); Okoth et al. (2014); Salvo and Valladares (1997); Shepard et al. (1998); Shiao and Wu (2000); Spencer (1973); Takano et al. (2008); Valladares et al. (1996); Valladares et al. (1999); Valladares et al. (2011); Weintraub et al. (2017); Yabas et al. (1995)
Commercial	Green pea	<i>Phaseolus vulgaris</i>	<i>vulgaris</i>	Fabaceae	Mujica and Kroschel (2011)	
Commercial	Garden pea; Pea; Snow pea; Sugar snap	<i>Pisum sativum</i>		Fabaceae	Sivapragasam et al. (1999); Foba et al. (2015); Larrain and Munoz (1997); Mujica and Kroschel (2011); Rauf et al. (2000); Scheffer et al. (2001); Sivapragasam and Syed (1999)	Andersen et al. (2002); Andersen et al. (2008); Bahlai et al. (2006); He et al. (2001); (Koch and Waterhouse 2000); Korytkowski (2014); Korytkowski (1982); Martin et al. (2005); Musundire et al. (2012); Okoth et al. (2014); Pang et al. (2006); Ripa et al. (1995); Salvo and Valladares (1997); Saunders et al. (1998); Silva (1993); Spencer (1973); Takano et al. (2008); Valladares et al. (1996); Valladares et al. (1999); Valladares et al. (2011); Wei et al. (2000); Weintraub et al. (2017); Yabas et al. (1995); Kox et al. (2005)
Commercial		<i>Pisum sativum</i>	<i>macrocarpenser</i>	Fabaceae		He et al. (2001)
Commercial	Snow pea	<i>Pisum sp.</i>		Fabaceae		Collins (1996); Scheffer et al. (2006); Shepard et al. (1998)
Commercial	Broad bean; Faba bean; Fava bean	<i>Vicia faba</i>		Fabaceae	Mujica and Kroschel (2011); Rauf et al. (2000); Salvo and Valladares (2002); Scheffer et al. (2001)	Blanchard (1938); Collins (1996); Echevarria et al. (1994); Gallegos 2000; Weintraub et al. (2017); He et al. (2001); He et al. (2002); (Koch and Waterhouse 2000); Korytkowski (1982); Musundire et al. 2011; Musundire et al. (2012); Noujeim et al. 2013; Okoth et al. (2014);

Host type	Common names	Scientific name	Variety	Family	Records with comprehensive evidence	All other records (partial and unverified records)
						Salvo and Valladares (1997); Saunders et al. (1998); Silva (1993); Takano et al. (2008); Valladares (1984); Valladares et al. (1996); Valladares et al. (1999); Valladares et al. (2011); Wei et al. (2000); Yabas et al. (1995)
Commercial		<i>Vicia</i> sp.		Fabaceae		Spencer (1973)
Commercial	Red bean; Black eyed pea; Cowpea	<i>Vigna sinensis</i>		Fabaceae	Rauf et al. (2000)	He et al. (2001); Shepard et al. (1998); Wei et al. (2000)
Commercial	Walp Cowpea; Yard-long bean; Long bean; Snakebean; Cowpea	<i>Vigna unguiculata</i>		Fabaceae	Foba et al. (2015); Rauf et al. (2000)	Hincapie et al. (1993)
Commercial		<i>Vigna unguiculata</i>	<i>unguiculata</i>	Fabaceae		Takano et al. (2008)
Commercial	Basil; Sweet basil; Thai basil	<i>Ocimum basilicum</i>		Lamiaceae	Mujica and Kroschel (2011); Rauf et al. (2000)	
Commercial	Okra	<i>Abelmoschus esculentus</i>		Malvaceae	Foba et al. (2015)	
Commercial	Pepper; Sweet pepper; Green pepper; Chili; Capsicum	<i>Capsicum annuum</i>		Solanaceae	Sivapragasam et al. (1999); Mujica and Kroschel (2011); Rauf et al. (2000); Sivapragasam and Syed (1999)	Calabretta et al. 1995; (Koch and Waterhouse 2000); Pang et al. (2006); Salvo and Valladares (1997); Saunders et al. (1998); Shepard et al. (1998); Shiao and Wu (2000); Spencer (1973); Valladares (1984); Valladares et al. (1996); Valladares et al. (1999); Valladares et al. (2011); Weintraub et al. (2017)

Host type	Common names	Scientific name	Variety	Family	Records with comprehensive evidence	All other records (partial and unverified records)
Commercial	Yellow chili	<i>Capsicum baccatum</i>		Solanaceae	Mujica and Kroschel (2011)	
Commercial	Sweet pepper; Pepper; Chilli	<i>Capsicum</i> sp.		Solanaceae	Foba et al. (2015)	Romero-Zuniga et al. (1991)
Commercial	Tomato	<i>Lycopersicon esculentum</i>		Solanaceae	Mujica and Kroschel (2011); Rauf et al. (2000); Sivapragasam et al. (1999); Sivapragasam and Syed (1999)	Korytkowski (2014); Kox et al. (2005); He et al. (2001); (Koch and Waterhouse 2000); Okoth et al. (2014); Olivera and Bordat (1996); Osorio et al. 1983; Salvo and Valladares (1997); Shepard et al. (1998); Spencer (1973); Valladares (1984); Valladares et al. (1996); Valladares et al. (1999); Valladares et al. (2011); Collins (1996);
Commercial	Tomato	<i>Lycopersicon esculentum</i>	<i>esculentum</i>	Solanaceae		Scheffer et al. (2006)
Commercial	Tomato	<i>Solanum lycopersicum</i>		Solanaceae	Foba et al. (2015)	Andersen et al. (2008); Andersen et al. (2002); de Clercq and Casteels (1992); Godinho and Mexia (2000); Hanafi (2005); Hidalgo and Carballo (1991); Korytkowski (1982); Macdonald et al. (2003); Moura et al. (2014); Musundire et al. (2012); Weintraub et al. (2017); Pang et al. (2006); Parish et al. (2017); Saunders et al. (1998); Weintraub et al. (2017)
Commercial	Brinjal eggplant;	<i>Solanum</i>		Solanaceae	Foba et al. (2015); Rauf et	He et al. (2001); Korytkowski (1982);

Host type	Common names	Scientific name	Variety	Family	Records with comprehensive evidence	All other records (partial and unverified records)
	Eggplant; common eggplant; Pickling eggplant	<i>melongena</i>			al. (2000)	Salvo and Valladares (1997); Saunders et al. (1998); Valladares (1984); Valladares et al. (1996); Valladares et al. (1999); Valladares et al. (2011)
Commercial		<i>Solanum sp.</i>		Solanaceae	Scheffer et al. (2001)	Andrade et al. (1989)
Commercial	Potato	<i>Solanum tuberosum</i>		Solanaceae	de Souza (1986); Foba et al. (2015); Larrain and Munoz (1997); Mujica and Kroschel (2011); Nino et al. (2009); Rauf et al. (2000); Salvo and Valladares (2002); Scheffer et al. (2001); Sivapragasam and Syed (1999)	Andersen et al. (2002); Andersen et al. (2008); Bahlai et al. (2006); Gallegos 2000; He et al. (2001); Hidalgo and Carballo (1991); Iwasaki (2004a); (Koch and Waterhouse 2000); Korytkowski (2014); Korytkowski (1982); Macdonald et al. (2003); Maharjan et al. 2014; Martin et al. (2005); Romero-Zuniga et al. (1991); Salvo and Valladares (1997); Saunders et al. (1998); Shepard et al. (1998); Spencer (1973); Valladares (1984); Valladares et al. (1996); Valladares et al. (1999); Valladares et al. (2011); Wijesekara (2010); Yabas et al. (1995)
Ornamental	Pot marigold	<i>Calendula officinalis</i>		Asteraceae	Salvo and Valladares (2002)	He et al. (2001); Korytkowski (1982); Salvo and Valladares (1997); Valladares (1984); Valladares et al. (1996); Valladares et al. (1999); Valladares et al. (2011)
Ornamental		<i>Calendula sp.</i>		Asteraceae		Weintraub et al. (2017)
Ornamental	Chinese aster	<i>Callistephus chinensis</i>		Asteraceae	Salvo and Valladares (2002)	de Clercq and Casteels (1992); Salvo and Valladares (1997); Valladares et al. (1999); Valladares et al. (2011); Wei et al. (2000); Valladares et al. (1996)
Ornamental	Garland chrysanthemum	<i>Chrysanthemum coronarium</i>		Asteraceae		Wei et al. (2000)

Host type	Common names	Scientific name	Variety	Family	Records with comprehensive evidence	All other records (partial and unverified records)
Ornamental	Garland chrysanthemum	<i>Chrysanthemum coronarium</i>	<i>coronarium</i>	Asteraceae	Tran (2009)	
Ornamental	Garland chrysanthemum; Chrysanthemum	<i>Chrysanthemum</i> sp.		Asteraceae	Scheffer et al. (2001)	Collins (1996); Hincapie et al. (1993); Pang et al. (2006); Salvo and Valladares (1997); Scheffer et al. (2006); Weintraub et al. (2017); Anderson et al. (2008)
Ornamental	Chrysanthemum	<i>Chrysanthemum</i> spp.		Asteraceae	Rauf et al. (2000)	
Ornamental	Dahlia	<i>Dahlia</i> sp.		Asteraceae	Rauf et al. (2000)	Collins (1996); Weintraub et al. (2017)
Ornamental	Gerbera; Barberton daisy; Transvaal daisy	<i>Gerbera jamesonii</i>		Asteraceae	Rauf et al. (2000)	Wei et al. (2000)
Ornamental		<i>Gerbera</i> sp.		Asteraceae		Weintraub et al. (2017)
Ornamental	Marigold; Aztec marigold	<i>Tagetes erecta</i>		Asteraceae	Mujica and Kroschel (2011)	Wei et al. (2000)
Ornamental		<i>Tagetes erecta</i>	<i>hybrids</i>	Asteraceae		de Clercq and Casteels (1992)
Ornamental	Marigold	<i>Tagetes</i> sp.		Asteraceae		Weintraub et al. (2017); Salvo and Valladares (1997); Valladares (1984); Valladares et al. (1996)
Ornamental		<i>Eustoma</i> sp.		Gentianaceae	Nakamura et al. (2013)	Weintraub et al. (2017)
Ornamental	'Swai tanah'	<i>Nasturtium indicum</i>		Tropaeolaceae	Rauf et al. (2000)	
Ornamental	Garden nasturtium	<i>Tropaeolum majus</i>		Tropaeolaceae	Salvo and Valladares (2002)	He et al. (2001); Salvo and Valladares (1997); Silva (1993); Valladares (1984)
Ornamental	Nasturtium	<i>Tropaeolum</i> sp.		Tropaeolaceae	Spencer (1983)	
Non cultivated	Amaranth; Bayam	<i>Amaranthus</i> sp.*		Amaranthaceae	Rauf et al. (2000); Sivapragasam et al. (1999)	Hidalgo and Carballo (1991); Romero-Zuniga et al. (1991); Shepard et al. (1998); Valladares et al. (1999); Valladares et al. (2011); Weintraub et al.

Host type	Common names	Scientific name	Variety	Family	Records with comprehensive evidence	All other records (partial and unverified records)
						(2017)
Non cultivated	Goosefoot; Lamb's quarters; Fat hen	<i>Chenopodium album</i> *		Amaranthaceae	Salvo and Valladares (2002)	He et al. (2001); Silva (1993)
Non cultivated	Amaranth	<i>Deeringia amaranthoides</i> *		Amaranthaceae	Rauf et al. (2000)	
Non cultivated	Penniwort	<i>Hydrocotyle</i> sp.*		Araliaceae	Salvo and Valladares (2002)	Salvo and Valladares (1997)
Non cultivated	Chick weed; Goatweed	<i>Ageratum conyzoides</i> *		Asteraceae	Rauf et al. (2000)	
Non cultivated	Common burdock	<i>Arctium minus</i> *		Asteraceae	Salvo and Valladares (2002)	Salvo and Valladares (1997); Valladares et al. (1999); Valladares et al. (2011)
Non cultivated	Sow thistle	<i>Emilia sonchifolia</i> *		Asteraceae	Rauf et al. (2000); Scheffer et al. (2001)	Hincapie et al. (1993); Scheffer et al. (2006)
Non cultivated	American burnweed; Pilewort; Fireweed	<i>Erechtites hieracifolia</i> *		Asteraceae	Rauf et al. (2000)	Romero-Zuniga et al. (1991)
Non cultivated	Indian lettuce	<i>Lactuca indica</i>		Asteraceae	Rauf et al. (2000); Sivapragasam et al. (1999)	
Non cultivated	Cinderella weed; Nodeweed	<i>Synedrella nodiflora</i> *		Asteraceae	Rauf et al. (2000)	
Non	Wild cucumber	<i>Melothria indica</i>		Cucurbitaceae	Rauf et al. (2000)	

Host type	Common names	Scientific name	Variety	Family	Records with comprehensive evidence	All other records (partial and unverified records)
cultivated						
Non cultivated	Cutleaf ground cherry; Mullaca; Wild gooseberry	<i>Physalis angulata*</i>		Solanaceae	Rauf et al. (2000)	Hidalgo and Carballo (1991)
Non cultivated	American black nightshade	<i>Solanum americanum</i>		Solanaceae	Rauf et al. (2000)	

Commercial, ornamental, and non-cultivated plants for which only partial evidence of lifecycle completion in the field for the serpentine leafminer (*Liriomyza huidobrensis*) has been provided within the scientific literature

This includes all records where:

- (1) Adults were reared from field collected plant material infested with larvae, but no subsequent identification via morphology or molecular diagnostics was reported; or
- (2) Laboratory feeding and oviposition preference experiments were conducted that did not confirm lifecycle completion; or
- (3) Identification via morphology or molecular diagnostics was reported, but collection methods in the field were not reported. Lifestyle completion cannot be confirmed in these instances due to the potential use of non-specific collection methods (e.g. sticky traps or sweep nets); or
- (4) Laboratory cultures confirmed lifecycle completion, but evidence of lifecycle completion in the field was not provided; or
- (5) Only common names of plant were given, but the scientific name could be confidently inferred; or
- (6) The record reports presence of *L. huidobrensis* on imported material detected at the border.

Records for plants identified only to the genus level (sp. or spp.) that did not meet the above conditions are included here only if there are other records for species within this genus that met one of these conditions outlined above. Records for specific varieties within a species that did not meet the above conditions are included only if there are other records for that same species that meet one of the conditions outlined above. Scientific names in the table appear as they were originally cited. A number of these names are now recognised as synonyms, however for brevity, we are not reporting the currently accepted names for these taxa (refer to WorldFloraOnline.org, Australian Plant Census, or Catalogue of Life for current taxonomic decisions). All common names associated with each scientific name, across all reports, are included. For all non-cultivated plants, and asterisk after the scientific name indicates that the Atlas of Living Australia reports this species present within Australia (Atlas of Living Australia 2020). Additionally, many plants labelled as commercial and ornamental have escaped cultivation and naturalised in Australia. Records that report comprehensive confirmations of lifecycle completion are included in Table 13, while unverified records are included in Table 15. *Liriomyza huidobrensis* is highly polyphagous and no guarantee can be made that this list is exhaustive.

Table 14 Commercial, ornamental and non-cultivated plants for which only partial evidence of lifecycle completion in the field for the serpentine leafminer (*Liriomyza huidobrensis*) has been provided within the scientific literature

Host type	Common names	Scientific name	Variety	Family	All records (including any unverified records)
Commercial	Quinoa	<i>Chenopodium quinoa</i>		Amaranthaceae	CABI (2019); Korytkowski (1982)
Commercial	Quinoa	<i>Chenopodium quinoa</i>	Amarilla de Maranganí	Amaranthaceae	Yabar et al. (2002)
Commercial	Quinoa	<i>Chenopodium quinoa</i>	Blanca de Junín	Amaranthaceae	Yabar et al. (2002)
Commercial		<i>Allium chinense</i>		Amaryllidaceae	He et al. (2001)
Commercial	Chive	<i>Allium schoenoprasum</i>		Amaryllidaceae	Spencer (1973)
Commercial		<i>Allium tuberosum</i>	<i>rottler</i>	Amaryllidaceae	He et al. (2001)
Commercial		<i>Bupleurum green</i> (most likely a <i>Bupleurum rotundifolium</i> cultivar)		Apiaceae	He et al. (2001)
Commercial		<i>Bupleurum</i> sp.		Apiaceae	Andersen and Hofsvang (2010)
Commercial	Coriander	<i>Coriandrum sativum</i>		Apiaceae	Weintraub et al. (2017); He et al. (2001); Koch and Waterhouse 2000
Commercial		<i>Daucus sativa</i>	<i>sativus</i>	Apiaceae	Wei et al. (2000); Hidalgo and Carballo (1991)
Commercial	Giant taro	<i>Alocasia macrorrhiza</i>		Araceae	Wei et al. (2000)
Commercial		<i>Colocasia esculenta</i>		Araceae	He et al. (2001)
Commercial	Carthamum; Safflower	<i>Carthamus tinctorius</i>		Asteraceae	He et al. (2001); Sunderlands et al. (1992)
Commercial		<i>Carthamus</i> sp.		Asteraceae	Andersen and Hofsvang (2010)

Host type	Common names	Scientific name	Variety	Family	All records (including any unverified records)
Commercial		<i>Cichorium endivia</i>		Asteraceae	de Goffau 1991
Commercial	Endive	<i>Cichorium</i> sp.		Asteraceae	Weintraub et al. (2017); Salvo and Valladares (1997)
Commercial		<i>Crassocephalum rubens</i>		Asteraceae	Shiao and Wu (2000)
Commercial	Artichoke	<i>Cynara cardunculus</i>	<i>scolymus</i>	Asteraceae	Larrain et al. 2013
Commercial		<i>Cynara scolymus</i>		Asteraceae	Korytkowski (2014)
Commercial		<i>Cynara scolymus</i>	<i>scolymus</i>	Asteraceae	Korytkowski (1982)
Commercial		<i>Basella rubra</i>		Basellaceae	He et al. (2001)
Commercial	Gai lan; Asian broccoli	<i>Brassica alboglabra</i>		Brassicaceae	Bahlai et al. (2006); Martin et al. (2005)
Commercial		<i>Brassica botrytis</i>		Brassicaceae	Wei et al. (2000)
Commercial		<i>Brassica campestris</i> spp.		Brassicaceae	He et al. (2001)
Commercial		<i>Brassica chinense</i>		Brassicaceae	Wei et al. (2000)
Commercial		<i>Brassica napus</i>		Brassicaceae	Wei et al. (2000)
Commercial		<i>Brassica pekinensis</i>		Brassicaceae	Wei et al. (2000)
Commercial		<i>Brassica pekinensis pekinensis</i>		Brassicaceae	He et al. (2002)
Commercial	Quinoa	<i>Chenopodium quinoa</i>		Chenopodiaceae	Korytkowski (2014)
Commercial		<i>Spinacea oleracea</i>		Chenopodiaceae	Korytkowski (2014)
Commercial		<i>Ipomoea aquatica</i>		Convolvulaceae	He et al. (2001)
Commercial	Wax gourd; White goard; Chinese melon; China squash; Ash gourd	<i>Benincasa hispida</i>		Cucurbitaceae	He et al. (2001)

Host type	Common names	Scientific name	Variety	Family	All records (including any unverified records)
Commercial	Watermelon	<i>Citrullus vulgaris</i>		Cucurbitaceae	Korytkowski (2014); Korytkowski (1982)
Commercial	Melon; Cantaloupe; Muskmelon; Honey melon; Cassaba melon	<i>Cucumis melo</i>		Cucurbitaceae	EPPO (2020); CABI (2019); Valladares et al. (2011); Guimarães et al. (2009); Koch and Waterhouse 2000; Valladares et al. (1999); Valladares et al. (1996); Valladares (1984); Spencer (1973);
Commercial	Bottle gourd	<i>Lagenaria siceraria</i>		Cucurbitaceae	Kox et al. (2005)
Commercial		<i>Lagenaria</i> sp.		Cucurbitaceae	Shiao and Wu (2000)
Commercial	Angled luffa; Sing kwa; Towelsponge; Angled luffa	<i>Luffa acutangula</i>		Cucurbitaceae	He et al. (2002)
Commercial	Towel gourd	<i>Luffa cylindrica</i>		Cucurbitaceae	Shiao and Wu (2000); Valladares et al. (1996)
Commercial	Creeper bean	<i>Dolichos lablab</i>		Fabaceae	He et al. (2001)
Commercial		<i>Lotus</i> sp.		Fabaceae	Collins (1996)
Commercial	White clover	<i>Trifolium repens</i>		Fabaceae	He et al. (2001); Wei et al. (2000); Andrade et al. (1989)
Commercial		<i>Trigonella</i> sp.		Fabaceae	Collins (1996)
Commercial		<i>Vicia fava</i>		Fabaceae	Korytkowski (2014)
Commercial		<i>Vicia sativa</i>		Fabaceae	Weintraub et al. (2017); He et al. (2001)
Commercial	Mungbean; Green gram	<i>Vigna radiata</i>		Fabaceae	Andersen et al. (2002)
Commercial	Spearmint	<i>Mentha spicata</i>		Lamiaceae	Wei et al. (2000)

Host type	Common names	Scientific name	Variety	Family	All records (including any unverified records)
Commercial		<i>Linum usitatissimum</i>		Linaceae	Spencer (1973)
Commercial	Barley	<i>Hordeum vulgare</i>		Poaceae	He et al. (2001)
Commercial		<i>Triticum aestivum</i>		Poaceae	He et al. (2001)
Commercial		<i>Zea mays</i>		Poaceae	He et al. (2001); Salvo and Valladares (1997)
Commercial	Wild rice	<i>Zizania latifolia</i>		Poaceae	Wei et al. (2000)
Commercial	Bell pepper	<i>Capsicum frutescens</i>		Solanaceae	He et al. (2002); He et al. (2001)
Commercial		<i>Lycium chinense</i>		Solanaceae	He et al. (2001)
Commercial		<i>Nicotiana</i> sp.		Solanaceae	Korytkowski (2014); Korytkowski (1982)
Commercial		<i>Nicotiana tabacum</i>		Solanaceae	Korytkowski (2014); He et al. (2001); Korytkowski (1982)
Commercial		<i>Solanum muricatum</i>		Solanaceae	Larrai 2002
Ornamental		<i>Thurnbergia alata</i>		Acanthaceae	Andersen and Hofsvang (2010)
Ornamental	Cock's comb	<i>Celosia argentea</i>		Amaranthaceae	He et al. (2001)
Ornamental		<i>Celosia argentea</i>	<i>cristata</i>	Amaranthaceae	He et al. (2001)
Ornamental		<i>Centella asiatica</i>		Apiaceae	He et al. (2001)
Ornamental		<i>Bupleurum griffithii</i>		Apiaceae	Andersen and Hofsvang (2010)
Ornamental		<i>Eryngium</i> sp.		Apiaceae	Andersen and Hofsvang (2010)
Ornamental		<i>Eryngium alpinum</i>		Apiaceae	Andersen and Hofsvang (2010)
Ornamental		<i>Eryngium foetidum</i>		Apiaceae	Andersen and Hofsvang (2010)
Ornamental		<i>Hydrocotyle umbellata</i>		Apiaceae	Korytkowski (2014)
Ornamental	Java waterdropwort	<i>Oenanthe javanica</i>		Apiaceae	He et al. (2001)

Host type	Common names	Scientific name	Variety	Family	All records (including any unverified records)
Ornamental		<i>Catharanthus roseus</i>		Apocynaceae	He et al. (2001)
Ornamental		<i>Chionodoxa luciliae</i>		Asparagaceae	He et al. (2001)
Ornamental		<i>Hemerocallis fulva</i>		Asphodelaceae	He et al. (2001)
Ornamental	Aster	<i>Aster</i> sp.		Asteraceae	Pang et al. (2006); Kox et al. (2005); Collins (1996)
Ornamental		<i>Argyranthemum frutescens</i>		Asteraceae	EPPO (2020); Andersen and Hofsvang (2010)
Ornamental		<i>Argyranthemum</i> sp.		Asteraceae	Weintraub et al. (2017); Andersen and Hofsvang (2010)
Ornamental	Daisy	<i>Bellis perennis</i>		Asteraceae	Valladares et al. (2011); He et al. (2001); Valladares et al. (1999); Salvo and Valladares (1997); Valladares (1984)
Ornamental		<i>Callendula officinallis</i>		Asteraceae	Korytkowski (2014)
Ornamental		<i>Centaurea cyanus</i>		Asteraceae	He et al. (2001)
Ornamental		<i>Centaurea</i> sp.		Asteraceae	Andersen and Hofsvang (2010)
Ornamental		<i>Chrysanthemum frutescens</i>		Asteraceae	de Clercq and Casteels (1992)
Ornamental	Oxeye daisy	<i>Chrysanthemum leucanthemum</i>		Asteraceae	Valladares et al. (2011); Valladares et al. (1999); Valladares et al. (1996); Valladares (1984);
Ornamental	Chrysanthemum; Florists' chrysanthemum	<i>Chrysanthemum morifolium</i>		Asteraceae	Valladares et al. (2011); Wei et al. (2000); Valladares et al. (1999); Valladares et al. (1996) de Clercq and Casteels (1992); Andrade et al. (1989); Valladares (1984)

Host type	Common names	Scientific name	Variety	Family	All records (including any unverified records)
Ornamental		<i>Cineraria cruenta</i>		Asteraceae	He et al. (2001)
Ornamental		<i>Cineraria</i> spp.		Asteraceae	Valladares (1984); Blanchard (1926)
Ornamental		<i>Conoclinium coelestinum</i>		Asteraceae	He et al. (2001)
Ornamental		<i>Cosmos bipinnatus</i>		Asteraceae	He et al. (2001)
Ornamental		<i>Craspedia globosa</i>		Asteraceae	He et al. (2001)
Ornamental	Dahlia	<i>Dahlia pinnata</i>		Asteraceae	He et al. (2001); Salvo and Valladares (1997); de Clercq and Casteels (1992);
Ornamental	Dahlia	<i>Dahlia variabilis</i>		Asteraceae	Weintraub et al. (2017); Valladares et al. (2011); Valladares et al. (1999); Valladares et al. (1996); Valladares (1984);
Ornamental		<i>Dendranthema morifolium</i>		Asteraceae	Cuthbertson et al. (2013)
Ornamental		<i>Gaillardia pulchella</i>		Asteraceae	He et al. (2001)
Ornamental		<i>Helianthus annuus</i>		Asteraceae	Korytkowski (2014)
Ornamental		<i>Helianthus</i> sp.		Asteraceae	Weintraub et al. (2017); Collins (1996);
Ornamental	Indian aster	<i>Kalimeris indica</i>		Asteraceae	Shiao and Wu (2000)
Ornamental		<i>Leucanthemum</i> sp.		Asteraceae	Andersen and Hofsvang (2010); Kox et al. (2005)
Ornamental		<i>Osteospermum</i> sp.		Asteraceae	Weintraub et al. (2017); Andersen and Hofsvang (2010)
Ornamental		<i>Pyrethrum cinerariifolium</i>		Asteraceae	He et al. (2001)

Host type	Common names	Scientific name	Variety	Family	All records (including any unverified records)
Ornamental	Goldenrod	<i>Solidago sp.</i>		Asteraceae	Andersen and Hofsvang (2010); He et al. (2001)
Ornamental	Marigold; French marigold	<i>Tagetes patula</i>		Asteraceae	Wei et al. (2000)
Ornamental		<i>Tagetes tenuitolia</i>		Asteraceae	Valladares et al. (2011); Valladares et al. (1999); Salvo and Valladares (1997); Valladares (1984);
Ornamental		<i>Tanacetum parthenium</i>		Asteraceae	EPPO (2020); He et al. (2001)
Ornamental		<i>Zinnia elegans</i>		Asteraceae	Valladares et al. (2011); He et al. (2001); Valladares et al. (1999); Valladares (1984)
Ornamental		<i>Zinnia augustifolia</i>		Asteraceae	Andersen and Hofsvang (2010)
Ornamental		<i>Zinnia sp.</i>		Asteraceae	Kox et al. (2005)
Ornamental		<i>Impatiens balsamina</i>		Balsaminaceae	He et al. (2001)
Ornamental		<i>Impatiens caeruleum</i>		Balsaminaceae	He et al. (2001)
Ornamental		<i>Eruca sp.</i>		Brassicaceae	Andersen and Hofsvang (2010)
Ornamental		<i>Eruca vesicaria</i>		Brassicaceae	Andersen and Hofsvang (2010)
Ornamental		<i>Lobularia maritima</i>		Brassicaceae	He et al. (2001)
Ornamental		<i>Matthiola incana</i>		Brassicaceae	He et al. (2001)
Ornamental		<i>Matthiola sp.</i>		Brassicaceae	He et al. (2001)
Ornamental		<i>Calceolaria crenatiflora</i>		Calceolariaceae	He et al. (2001)
Ornamental		<i>Campanula medium</i>		Campanulaceae	He et al. (2001)
Ornamental		<i>Platycodon grandifloras</i>		Campanulaceae	He et al. (2001)

Host type	Common names	Scientific name	Variety	Family	All records (including any unverified records)
Ornamental		<i>Trachelium</i> sp.		Campanulaceae	Andersen and Hofsvang (2010)
Ornamental		<i>Dianthus barbatus</i>		Caryophyllaceae	He et al. (2001)
Ornamental	Carnation	<i>Dianthus caryophyllus</i>		Caryophyllaceae	EPPO (2020); Valladares et al. (2011); He et al. (2001); (Koch and Waterhouse 2000); Valladares et al. (1999); Shepard et al. (1998); Salvo and Valladares (1997); Valladares (1984)
Ornamental	Indian pink; China pink	<i>Dianthus chinensis</i>		Caryophyllaceae	He et al. (2001); de Clercq and Casteels (1992)
Ornamental		<i>Dianthus hybridus</i>		Caryophyllaceae	He et al. (2001)
Ornamental		<i>Dianthus</i> sp.		Caryophyllaceae	Andersen and Hofsvang (2010); Kox et al. (2005); Collins (1996);
Ornamental	Showy baby's breath	<i>Gypsophila elegans</i>		Caryophyllaceae	Valladares et al. (2011); He et al. (2002); He et al. (2001); Valladares et al. (1999)
Ornamental		<i>Gypsophila muralis</i>		Caryophyllaceae	Kox et al. (2005)
Ornamental	Gypsophila; Baby's breath	<i>Gypsophila paniculata</i>		Caryophyllaceae	Cure and Cantor 2003; He et al. (2001); Wei et al. (2000)
Ornamental	Gypsophila	<i>Gypsophila</i> sp.		Caryophyllaceae	Weintraub et al. (2017); Collins (1996); Andrade et al. (1989); Sunderlands et al. (1992)
Ornamental		<i>Gypsophila perfecta</i>		Caryophyllaceae	Andersen and Hofsvang (2010)
Ornamental		<i>Pharbitis purpurea</i>		Convolvulaceae	He et al. (2001)

Host type	Common names	Scientific name	Variety	Family	All records (including any unverified records)
Ornamental		<i>Euphorbia marginata</i>		Euphorbiaceae	He et al. (2001)
Ornamental		<i>Lathyrus latifolius</i>		Fabaceae	Valladares et al. (2011); Valladares et al. (1999); Valladares (1984)
Ornamental		<i>Lathyrus odoratus</i>		Fabaceae	Valladares et al. (2011); He et al. (2001); Valladares et al. (1999); Valladares (1984)
Ornamental		<i>Lupin</i> sp.		Fabaceae	Collins (1996)
Ornamental		<i>Lupinus rassel</i>	<i>Russell hybrid</i>	Fabaceae	He et al. (2001)
Ornamental		<i>Lupinus</i> sp.		Fabaceae	Kox et al. (2005)
Ornamental	Fenugreek	<i>Trigonella foenum-graecum</i>		Fabaceae	Andersen and Hofsvang (2010)
Ornamental	Exacum	<i>Exacum</i> sp.		Gentianaceae	Andersen and Hofsvang (2010)
Ornamental		<i>Exacum affine</i>		Gentianaceae	Andersen and Hofsvang (2010)
Ornamental		<i>Eustoma grandiflorum</i>		Gentianaceae	Andersen and Hofsvang (2010)
Ornamental		<i>Eustoma russellianum</i>		Gentianaceae	He et al. (2001); Wei et al. (2000)
Ornamental		<i>Lisianthus</i> sp.		Gentianaceae	Andersen and Hofsvang (2010); Weintraub et al. (2017); Collins (1996)
Ornamental		<i>Lisianthus russelianus</i>		Gentianaceae	Andersen and Hofsvang (2010)
Ornamental		<i>Scaevola aemula</i>		Goodeniaceae	Andersen and Hofsvang (2010)
Ornamental		<i>Hydrangea macrophylla</i>		Hydrangeaceae	He et al. (2001)
Ornamental		<i>Freesia refracta</i>		Iridaceae	He et al. (2001)
Ornamental	Sword lily	<i>Gladiolus hybridus</i>		Iridaceae	He et al. (2001); Wei et al. (2000)
Ornamental		<i>Lamium</i> sp.		Lamiaceae	Andersen and Hofsvang (2010)

Host type	Common names	Scientific name	Variety	Family	All records (including any unverified records)
Ornamental		<i>Moluccella laevis</i>		Lamiaceae	He et al. (2001)
Ornamental		<i>Moluccella</i> sp.		Lamiaceae	Andersen and Hofsvang (2010)
Ornamental		<i>Salvia splendens</i>		Lamiaceae	He et al. (2001)
Ornamental		<i>Lilium davidii</i>		Liliaceae	He et al. (2001)
Ornamental		<i>Lilium longiflorum</i>		Liliaceae	He et al. (2001)
Ornamental		<i>Torenia</i> sp.		Linderniaceae	Kox et al. (2005)
Ornamental		<i>Althaea rosea</i>		Malvaceae	Korytkowski (2014)
Ornamental		<i>Hibiscus trionum</i>		Malvaceae	He et al. (2001)
Ornamental		<i>Godetia amoena</i>		Onagraceae	He et al. (2001)
Ornamental		<i>Oenothera rosea</i>		Onagroideae	He et al. (2001)
Ornamental		<i>Papaver rhoeas</i>		Papaveraceae	Valladares et al. (2011); He et al. (2001); Valladares et al. (1999); Salvo and Valladares (1997); Valladares (1984)
Ornamental	Snapdragon	<i>Antirrhinum majus</i>		Plantaginaceae	Martin et al. (2005)
Ornamental		<i>Limonium latifolium</i>		Plumbaginaceae	He et al. (2001)
Ornamental		<i>Limonium tataricum</i>		Plumbaginaceae	He et al. (2001)
Ornamental		<i>Myosotis sylvatica</i>		Plumbaginaceae	He et al. (2001)
Ornamental		<i>Lagurus ovatus</i>		Poaceae	He et al. (2001)
Ornamental	Phlox	<i>Phlox drummondii</i>		Polemoniaceae	Valladares et al. (2011); He et al. (2001); Valladares et al. (1999); Salvo and Valladares (1997); de Clercq and Casteels (1992); Valladares (1984)
Ornamental		<i>Primula acaulis</i>		Primulaceae	He et al. (2001)

Host type	Common names	Scientific name	Variety	Family	All records (including any unverified records)
Ornamental		<i>Primula obconica</i>		Primulaceae	He et al. (2001); Salvo and Valladares (1997); de Clercq and Casteels (1992)
Ornamental		<i>Delphinium grandiflorum</i>		Ranunculaceae	He et al. (2001)
Ornamental		<i>Delphinium</i> sp.		Ranunculaceae	He et al. (2001)
Ornamental		<i>Nigella damascena</i>		Ranunculaceae	He et al. (2001)
Ornamental		<i>Ranunculus asiaticus</i>		Ranunculaceae	Valladares et al. (2011); He et al. (2001); Valladares et al. (1999); Valladares et al. (1996)
Ornamental		<i>Ranunculus sceleratus</i>		Ranunculaceae	Wei et al. (2000)
Ornamental	Buttercup	<i>Ranunculus</i> sp.		Ranunculaceae	Collins (1996)
Ornamental		<i>Ranunculus viridis</i>		Ranunculaceae	He et al. (2001)
Ornamental		<i>Rosa chinese</i>		Rosaceae	Wei et al. (2000)
Ornamental		<i>Rosa</i> sp.		Rosaceae	Andersen and Hofsvang (2010)
Ornamental		<i>Diascia</i> sp.		Scrophulariaceae	Andersen and Hofsvang (2010); Saethre (1996)
Ornamental		<i>Nicotiana alata</i>		Solanaceae	de Clercq and Casteels (1992)
Ornamental		<i>Petunia hybrida</i>		Solanaceae	He et al. (2001)
Ornamental	Petunia	<i>Petunia</i> sp.		Solanaceae	Weintraub et al. (2017); Valladares et al. (2011); Valladares et al. (1999); Salvo and Valladares (1997); Collins (1996); Hincapie et al. (1993)
Ornamental		<i>Petunia</i> spp.		Solanaceae	Valladares (1984)
Ornamental		<i>Petunia</i> spp.	<i>hybrids</i>	Solanaceae	de Clercq and Casteels (1992)
Ornamental	Vervain	<i>Verbena officinalis</i>		Verbenaceae	He et al. (2001)

Host type	Common names	Scientific name	Variety	Family	All records (including any unverified records)
Ornamental		<i>Verbena</i> sp.		Verbenaceae	Weintraub et al. (2017); Collins (1996); de Clercq and Casteels (1992)
Ornamental		<i>Viola tricolor</i>		Violaceae	Valladares et al. (2011); Valladares et al. (1999); Salvo and Valladares (1997); Valladares (1984)
Ornamental		<i>Viola tricolor</i>	<i>vannortensis</i>	Violaceae	He et al. (2001)
Ornamental		<i>Viola yedoensis</i>		Violaceae	He et al. (2001)
Non cultivated		<i>Sagittaria sagittifolia</i> *		Alismataceae	He et al. (2001)
Non cultivated		<i>Alstroemeria</i> sp.*		Alstroemeriaceae	Collins (1996)
Non cultivated		<i>Alstroemeria aurea</i> *		Alstroemeriaceae	He et al. (2001)
Non cultivated		<i>Amaranthus caudatus</i> *		Amaranthaceae	He et al. (2001)
Non cultivated		<i>Amaranthus lividus</i> *		Amaranthaceae	He et al. (2001)
Non cultivated		<i>Amaranthus manostanus</i>		Amaranthaceae	He et al. (2001)
Non cultivated	Rough pigweed	<i>Amaranthus retroflexus</i> *		Amaranthaceae	He et al. (2001); Hincapie et al. (1993)
Non cultivated		<i>Amaranthus viridis</i> *		Amaranthaceae	Shiao and Wu (2000); Verjel-Manzano and Mejia-Florez (2000); Spencer (1973)
Non cultivated		<i>Chenopodium ambrosioides</i> * (Korytkowski (2014) reports as <i>C. abrossoides</i>)		Amaranthaceae	Korytkowski (2014); Valladares et al. (2011); He et al. (2001); Valladares et al. (1999); Valladares (1984); Korytkowski (1982)
Non cultivated		<i>Chenopodium abrossoides</i>		Amaranthaceae	
Non cultivated		<i>Chenopodium hirsutum</i>		Amaranthaceae	Korytkowski (2014)

Host type	Common names	Scientific name	Variety	Family	All records (including any unverified records)
Non cultivated		<i>Chenopodium murale</i> *		Amaranthaceae	Korytkowski (1982)
Non cultivated		<i>Chenopodium paniculatum</i>		Amaranthaceae	Andrade et al. (1989)
Non cultivated		<i>Gomphrena globosa</i> *		Amaranthaceae	He et al. (2001)
Non cultivated		<i>Iresine herbstii</i> *		Amaranthaceae	Wei et al. (2000)
Non cultivated		<i>Unknown Amaranthaceae</i>		Amaranthaceae	Salvo and Valladares (1997)
Non cultivated		<i>Hydrocotyle umbellata</i> *		Araliaceae	Hincapie et al. (1993); Hidalgo and Carballo (1991); Korytkowski (1982)
Non cultivated		<i>Arctium lappa</i> *		Asteraceae	Scheffer et al. (2006); Silva (1993)
Non cultivated		<i>Artemisia annua</i>		Asteraceae	He et al. (2001)
Non cultivated	Black jack; Spanish needle; hairy beggarticks	<i>Bidens pilosa</i> *		Asteraceae	Weintraub et al. (2017); Pang et al. (2006); Hidalgo and Carballo (1991); Romero-Zuniga et al. (1991)
Non cultivated	Bidens	<i>Bidens sp.</i> *		Asteraceae	Salvo and Valladares (1997)
Non cultivated		<i>Carduus crispus</i>		Asteraceae	He et al. (2001)
Non cultivated		<i>Conyza canadensis</i> *		Asteraceae	Wei et al. (2000)
Non cultivated		<i>Conyza sp.</i> *		Asteraceae	Valladares et al. (2011); Valladares et al. (1999); Salvo and Valladares (1997)
Non cultivated		<i>Crepis pulchra</i>		Asteraceae	Salvo and Valladares (1997)
Non cultivated		<i>Dichrocephala auriculata</i>		Asteraceae	He et al. (2001)
Non cultivated		<i>Echinops ritro</i>		Asteraceae	He et al. (2001)
Non cultivated		<i>Erigeron breviscapus</i>		Asteraceae	He et al. (2001)
Non cultivated		<i>Galinsoga caracasana</i>		Asteraceae	Andrade et al. (1989); Spencer (1983); Spencer (1973)

Host type	Common names	Scientific name	Variety	Family	All records (including any unverified records)
Non cultivated	Quickweed; Shaggy soldier	<i>Galinsoga ciliata</i>		Asteraceae	Hidalgo and Carballo (1991); Romero-Zuniga et al. (1991); Andrade et al. (1989)
Non cultivated		<i>Galinsoga parviflora</i> *		Asteraceae	Wei et al. (2000); Hincapie et al. (1993)
Non cultivated		<i>Helichrysum bracteatum</i> *		Asteraceae	Wei et al. (2000)
Non cultivated		<i>Helipterum roseum</i> *		Asteraceae	He et al. (2001)
Non cultivated		<i>Hemistepta lyrata</i> *		Asteraceae	He et al. (2001)
Non cultivated	Spiny sow thistle; Prickly sow thistle	<i>Sonchus asper</i> *		Asteraceae	He et al. (2001); Hincapie et al. (1993)
Non cultivated		<i>Sonchus brachyotus</i>		Asteraceae	He et al. (2001)
Non cultivated		<i>Sonchus lunatus</i>		Asteraceae	Andrade et al. (1989)
Non cultivated	Common sow thistle	<i>Sonchus oleraceus</i> *		Asteraceae	Weintraub et al. (2017); He et al. (2001); Hidalgo and Carballo (1991); Romero-Zuniga et al. (1991)
Non cultivated		<i>Sonchus sp.</i> *		Asteraceae	Korytkowski (2014); Korytkowski (1982)
Non cultivated		<i>Taraxacum mongolicum</i>		Asteraceae	He et al. (2001)
Non cultivated		<i>Barbarea sp.</i> *		Brassicaceae	Shiao and Wu (2000)
Non cultivated		<i>Capsella bursa-pastoris</i> *		Brassicaceae	Weintraub et al. (2017); He et al. (2001); Andrade et al. (1989)
Non cultivated		<i>Cardamine hirsute</i> *		Brassicaceae	He et al. (2001)
Non cultivated		<i>Rorippa indica</i>		Brassicaceae	Scheffer et al. (2006); He et al. (2001)
Non cultivated		<i>Rorippa montana</i>		Brassicaceae	He et al. (2001)
Non cultivated		<i>Rorippa palustris</i> *		Brassicaceae	He et al. (2001)

Host type	Common names	Scientific name	Variety	Family	All records (including any unverified records)
Non cultivated		<i>Malachium aguaticum</i>		Caryophyllaceae	He et al. (2001)
Non cultivated		<i>Silene gallica</i> *		Caryophyllaceae	Andrade et al. (1989)
Non cultivated		<i>Stellaria media</i> *		Caryophyllaceae	Mujica (2007); He et al. (2001); Wei et al. (2000)
Non cultivated		<i>Stellaria yunnansis</i>		Caryophyllaceae	He et al. (2001)
Non cultivated		<i>Calystegia hederacea</i>		Convolvulaceae	He et al. (2001)
Non cultivated		<i>Calystegia sepium</i> *		Convolvulaceae	Wei et al. (2000)
Non cultivated	Castor bean; Castor oil bush	<i>Ricinus communis</i> *		Euphorbiaceae	He et al. (2001)
Non cultivated	Black medic	<i>Medicago lupulina</i> *		Fabaceae	He et al. (2001)
Non cultivated		<i>Medicago minima</i> *		Fabaceae	He et al. (2001)
Non cultivated		<i>Melilotus suaveolens</i>		Fabaceae	He et al. (2001)
Non cultivated		<i>Vicia tetrasperma</i> *		Fabaceae	He et al. (2001)
Non cultivated		<i>Leonurus heterophyllus</i>		Lamiaceae	He et al. (2001)
Non cultivated		<i>Althaea rosea</i>		Malvaceae	He et al. (2001); Korytkowski (1982)
Non cultivated		<i>Malva rotundifolia</i>		Malvaceae	He et al. (2001)
Non cultivated		<i>Malva verticillate</i> *		Malvaceae	He et al. (2001); Wei et al. (2000)
Non cultivated		<i>Sida</i> sp.*		Malvaceae	Scheffer et al. (2006)
Non cultivated		<i>Stephania delavayi</i>		Menispermaceae	He et al. (2001)
Non cultivated		<i>Oxalis corniculata</i> *		Oxalidaceae	He et al. (2001)
Non cultivated		<i>Plantago asiatica</i>		Plantaginaceae	He et al. (2001)
Non cultivated	Broad leaved plantain	<i>Plantago major</i> *		Plantaginaceae	He et al. (2001)
Non cultivated		<i>Veronica anagallis-aquatica</i> *		Plantaginaceae	He et al. (2001)

Host type	Common names	Scientific name	Variety	Family	All records (including any unverified records)
Non cultivated		<i>Setaria viridis</i> *		Poaceae	Wei et al. (2000)
Non cultivated		<i>Polygonum amphibium</i>		Polygonaceae	He et al. (2001)
Non cultivated	Knotweed	<i>Polygonum aviculare</i> *		Polygonaceae	He et al. (2001)
Non cultivated		<i>Polygonum hydropiper</i> *		Polygonaceae	He et al. (2001)
Non cultivated		<i>Polygonum nepalense</i>		Polygonaceae	He et al. (2001)
Non cultivated		<i>Rumex acetosa</i> *		Polygonaceae	He et al. (2001)
Non cultivated		<i>Nemesia strumose</i> *		Scrophulariaceae	He et al. (2001)
Non cultivated		<i>Datura ferox</i> *		Solanaceae	Valladares et al. (2011); Valladares et al. (1999); Salvo and Valladares (1997)
Non cultivated		<i>Datura stramonium</i> *		Solanaceae	Mujica (2007); He et al. (2001); Hincapie et al. (1993)
Non cultivated		<i>Nicotiana glauca</i> *		Solanaceae	Korytkowski (2014); Korytkowski (1982)
Non cultivated		<i>Nicotiana physaloides</i>		Solanaceae	He et al. (2001)
Non cultivated	Black nightshade	<i>Solanum nigrum</i> *		Solanaceae	He et al. (2001)
Non cultivated		<i>Solanum oleracelus</i>		Solanaceae	Noujeim et al. (2013)

Commercial, ornamental and non-cultivated plants with unverified records of lifecycle completion in the field for the serpentine leafminer (*Liriomyza huidobrensis*)

This includes all records for which no evidence was presented within the record for lifecycle completion and no evidence was shown for confident species identification, or evidence could not suitably be accessed for verification. We are not disputing the veracity of these records - we simply did not have enough information available to verify if these hosts support lifecycle completion in the field. This includes all records where:

- (1) Adults were collected by sweep netting the presumed host, but neither larval activity nor lifecycle completion was confirmed; or
- (2) A scientific name could not reliably be assumed from a reported common name; or
- (3) Collection and identification methodology were unclear or unreported; or
- (4) The record could not be accessed or suitably translated to confirm collection and identification methodology.

For some unverified species within this table, there may be other congeneric species that have been either comprehensively or partially confirmed as a host plant, and thus these would appear within Table 13 and Table 14. Scientific names in the table appear as they were originally cited. A number of these names are now recognised as synonyms, however for brevity, we are not reporting the currently accepted names for these taxa (refer to WorldFloraOnline.org, Australian Plant Census, or Catalogue of Life for current taxonomic decisions). For all non-cultivated plants, and asterisk after the scientific name indicates that the Atlas of Living Australia reports this species present within Australia (Atlas of Living Australia 2020). All common names associated with each scientific name, across all reports, are included. Additionally, many plants labelled as commercial and ornamental have escaped cultivation and naturalised in Australia. Records that report comprehensive confirmations of lifecycle completion are included in Table 13, while records that report only partial confirmations of lifecycle completion are included in Table 14. *Liriomyza huidobrensis* is highly polyphagous and no guarantee can be made that this list is exhaustive.

Table 15 Commercial, ornamental and non-cultivated plants with unverified records of lifecycle completion in the field for the serpentine leafminer (*Liriomyza huidobrensis*).

Host type	Common names	Scientific name	Variety	Family	Records
Commercial		<i>Sambucus</i> sp.		Adoxaceae	Weintraub et al. (2017)
Commercial	Leek	<i>Allium porrum</i>		Amaryllidaceae	Weintraub et al. (2017)
Commercial		<i>Levisticum officinale</i>		Apiaceae	Weintraub et al. (2017)
Commercial		<i>Petroselinum</i> sp.		Apiaceae	Hincapie et al. (1993)
Commercial		<i>Asparagus officinalis</i>		Asparagaceae	Silva (1993)

Host type	Common names	Scientific name	Variety	Family	Records
Commercial		<i>Cichorium intybus</i>		Asteraceae	Valladares et al. (1996); Valladares (1984); Valladares et al. (1999); Valladares et al. (2011); Weintraub et al. (2017)
Commercial		<i>Cannabis sativa</i>		Cannabaceae	EPPO (2020)
Commercial		<i>Humulus scandens</i>		Cannabaceae	Weintraub et al. (2017)
Commercial	Common cornsalad	<i>Valerianella locusta</i>		Caprifoliaceae	CABI (2019)
Commercial	Lentil; Lentil bean	<i>Lens culinaris</i>		Fabaceae	(Koch and Waterhouse 2000)
Commercial	Black gram	<i>Vigna mungo</i>		Fabaceae	Weintraub et al. (2017)
Commercial	Passionfruit	<i>Passiflora</i> sp.		Passifloraceae	Kahinga et al. (2017)
Ornamental		<i>Oenanthе benghalensis</i>		Apiaceae	Weintraub et al. (2017)
Ornamental		<i>Chrysanthemum indicum</i>		Asteraceae	EPPO (2020)
Ornamental		<i>Dahlia imperialis</i>		Asteraceae	Spencer (1983)
Ornamental		<i>Dendranthema x grandiflorum</i>		Asteraceae	EPPO (2020)
Ornamental	Sunflower; Common sunflower	<i>Helianthus annuus</i>		Asteraceae	Korytkowski (1982); Valladares et al. (1999); Valladares et al. (2011)
Ornamental		<i>Leucanthemum vulgare</i>		Asteraceae	EPPO (2020)
Ornamental		<i>Leucanthemum x superbum</i>		Asteraceae	EPPO (2020)
Ornamental	Butterweed	<i>Senecio cruentus</i>		Asteraceae	Calabretta et al. 1995
Ornamental		<i>Tanacetum vulgare</i>		Asteraceae	EPPO (2020)
Ornamental		<i>Primula x polyanthus</i>		Primulaceae	EPPO (2020)
Ornamental		<i>Ranunculus chinensis</i>		Ranunculaceae	Weintraub et al. (2017)
Ornamental		<i>Ranunculus sieboldii</i>		Ranunculaceae	Weintraub et al. (2017)
Ornamental		<i>Viola philippica</i>		Violaceae	Weintraub et al. (2017)
Non cultivated		<i>Trianthema portulacastrum</i> *		Aizoaceae	Mujica (2007)

Host type	Common names	Scientific name	Variety	Family	Records
Non cultivated		<i>Alternanthera philox</i>	<i>eroides</i>	Amaranthaceae	Weintraub et al. (2017)
Non cultivated		<i>Amaranthus hybridus</i> *		Amaranthaceae	Mujica (2007)
Non cultivated		<i>Chenopodium hircinum</i>		Amaranthaceae	Korytkowski (1982)
Non cultivated		<i>Artemisia argyi</i>		Asteraceae	Weintraub et al. (2017)
Non cultivated		<i>Carduus nutans</i> *		Asteraceae	Weintraub et al. (2017)
Non cultivated		<i>Conyza bonariensis</i> *		Asteraceae	Weintraub et al. (2017)
Non cultivated		<i>Eclipta prostrata</i> *		Asteraceae	Weintraub et al. (2017)
Non cultivated		<i>Galinsoga urticifolia</i>		Asteraceae	Weintraub et al. (2017)
Non cultivated		<i>Gnaphalium affine</i>		Asteraceae	Weintraub et al. (2017)
Non cultivated		<i>Gynura crepidioides</i> *		Asteraceae	Weintraub et al. (2017)
Non cultivated		<i>Schistocarpa platyphylla</i>		Asteraceae	Weintraub et al. (2017)
Non cultivated		<i>Taraxacum officinale</i> *		Asteraceae	Valladares et al. (1999); Valladares et al. (2011)
Non cultivated		<i>Diplotaxis muralis</i> *		Brassicaceae	Mujica (2007)
Non cultivated		<i>Stellaria alsine</i>		Caryophyllaceae	Weintraub et al. (2017)
Non cultivated		<i>Vaccaria pyramidata</i> *		Caryophyllaceae	Iwasaki (2004a)
Non cultivated	Chipilín	<i>Crotalaria longirostrata</i>		Fabaceae	Weintraub et al. (2017)
Non cultivated		<i>Streptocarpus sp.</i>		Gesneriaceae	Weintraub et al. (2017)
Non cultivated		<i>Leonurus sibiricus</i>		Lamiaceae	Weintraub et al. (2017)
Non cultivated		<i>Stachys arvensis</i> *		Lamiaceae	Mujica (2007)
Non cultivated		<i>Alcea sp.</i> *		Malvaceae	Weintraub et al. (2017)
Non cultivated	Purslane	<i>Portulaca oleracea</i> *		Portulacaceae	CABI (2019); Hincapie et al. (1993)

References

- Andersen A, Hofsvang T (2010) Pest risk assessment of the South American Leafminer (*Liriomyza huidobrensis*) in Norway. VKM, Oslo, Norway
- Andersen A, Nordhus E, Thang VT, et al (2002) Polyphagous *Liriomyza* species (Diptera: Agromyzidae) vegetables in Vietnam. 79:241–246
- Andersen A, Tran TTA, Nordhus E (2008) Distribution and importance of polyphagous *Liriomyza* species (Diptera, Agromyzidae) in vegetables in Vietnam. Nor J Entomol 55:149–164
- Andrade ME, Briceño JA, de Hoyos PM, Jiménez J (1989) Busqueda y reconocimiento de los enemigos naturales y hospedantes alternos de las principales plagas. en flores bajo invernadero en la sabana de bogota. Acta Biol Colomb 1:45–57
- Atlas of Living Australia website at <http://www.ala.org.au>. Accessed 28 January 2020.
- Ávalos DS, Ricobelli G, Palacios SM, Defagó MT (2013) Evaluación de la preferencia de *Liriomyza* spp. En genotipos diferentes de garbanzo y efecto del extracto de *Melia azedarach* L.: Resultados preliminares. Rev la Fac Ciencias Agrar 45:65–73
- Bahlai CA, Goodfellow SA, Stanley-horn DE, Hallett RH (2006) Endoparasitoid assemblage of the pea leafminer, *Liriomyza huidobrensis* (Diptera: Agromyzidae), in Southern Ontario. Environ Entomol 35:351–357. <https://doi.org/10.1603/0046-225x-35.2.351>
- Blanchard EE (1926) A dipterous leaf-miner on Cineraria, new to Science. Rev la Soc Entomológica Argentina 1:10–11
- Blanchard EE, Pisa LW, Amaral-Rogers V, et al (1938) Descripciones y anotaciones de dipteros argentinos. In: Anales de la Sociedad Científica Argentina. pp 345–386
- CABI (2019) *Liriomyza huidobrensis* (serpentine leafminer). In: Invasive Species Compend. www.cabi.org/isc/datasheet/30956
- Calabretta C, Calabro M, Colombo A, Campo G (1995) [Spreading of *Liriomyza huidobrensis* (Blanchard) (Diptera, Agromyzidae) in protected cultures in Sicily]. [Italian]. Informatore Fitopatologico.
- Carballo, M. R, Leon G, Ramirez A (1990) Biological control of *Liriomyza* sp. (Diptera: Agromyzidae) in horticultural crops of Costa Rica. Manejo Integr Plagas 16:4–11
- Clercq RD, Casteels H (1992) Occurrence and host plants of *Frankliniella occidentalis* and *Liriomyza huidobrensis* in Belgium. OEPP/EPPO Bull 22:363–366
- Collins DW (1996) The separation of *Liriomyza huidobrensis* (Diptera: Agromyzidae) from related indigenous and non-indigenous species encountered in the United Kingdom using cellulose acetate electrophoresis. Ann Appl Biol 128:387–398. <https://doi.org/10.1111/j.1744-7348.1996.tb07100.x>
- Cure JR, Cantor F (2003) Atividade predadora e parasítica de *Diglyphus begini* (Ashm.) (Hymenoptera: Eulophidae) sobre *Liriomyza huidobrensis* (Blanch.) (Diptera: Agromyzidae) em Cultivos de *Gypsophila paniculata* L. Neotrop Entomol 32:85–89. <https://doi.org/10.1590/S1519-566X2003000100012>
- Cuthbertson AGS, Mills K, Wonter-Smith T, et al (2013) Environmental evaluation of alternative chemicals to methyl bromide for fumigation of quarantine pests in transit. Int J Environ Sci Technol 10:1057–1066. <https://doi.org/10.1007/s13762-013-0237-9>
- De Goffau LJW (1991) *Liriomyza huidobrensis* (Blanchard) (Diptera:Agromyzidae) A new economically important leaf miner in the Netherlands. Proc Sect Exp Appl Entomol Netherlands Entomol Soc 2:41–45
- de Souza JC (1986) Levantamento das especies de moscas-minadoras (Diptera: Agromyzidae) em algumas culturas no estado de minas gerias. An da Soc Entomol do Bras 15:171–172
- Echevarría A, Gimeno C, Jimenez Ródena J (1994) *Liriomyza huidobrensis* (Blanchard, 1926) (Diptera, Agromyzidae) una nueva plaga en cultivos valencianos. Boletín Sanid Veg Plagas 20:103–109
- EPPO (2020) *Liriomyza huidobrensis*. In: EPPO Glob. Database. <https://gd.eppo.int/taxon/LIRIHU>. Accessed 12 Nov 2020
- Foba CN, Salifu D, Lagat ZO, et al (2015) Species composition, distribution, and seasonal abundance of

- Liriomyza* leafminers (Diptera: Agromyzidae) under Different Vegetable Production Systems and Agroecological Zones in Kenya. *Environ Entomol* 44:223–232. <https://doi.org/10.1093/ee/nvu065>
- Gallegos P (2000) Situación actual y perspectivas de control del “minador de la hoja” *Liriomyza huidobrensis*, en el cultivo de papa, en la provincia del Carchi
- Gloria R, Salas VH (1976) Control Químico de la “Mosca Minadora” *Liriomyza huidobrensis* Blanchard en Alfalfa. *Rev Peru Entomol* 19:89–91
- Godinho M, Mexia A (2000) Leafminers (*Liriomyza* sp.) importance in greenhouses in the Oeste region of Portugal and its natural parasitoids as control agents in IPM programs. 23:157–161
- Guimarães JA, Filho MM, Oliveira VR, et al (2009) Biología e manejo de mosca minadora no meloeiro. 1–9
- Hanafi A (2005) Invasive pests and diseases: A challenge to IPM in greenhouse crops. *Phytoparasitica* 33:423–426. <https://doi.org/10.1007/BF02981390>
- He C, Wu W, Wang S, Wang L (2001) Host plants and feeding preference of *Liriomyza huidobrensis*. *Acta Entomol Sin* 44:384–388
- He L, Zhang Y, Xiao N, et al (2002) *Liriomyza huidobrensis* in Yunnan, China: Current distribution and genetic structure of a recently established population. *Entomol Exp Appl* 102:213–219. <https://doi.org/10.1023/A:1019578824443>
- Hidalgo JE, Carballo VM (1991) Influencia de las malezas sobre los insectos controladores naturales de *Liriomyza huidobrensis* (Blanchard), (Diptera: Agromyzidae). (The influence of weeds on the natural enemies of *Liriomyza huidobrensis* (Blanchard) (Diptera:Agromyzidae)). *Manejo Integr Plagas* 20–21:49–54
- Hincapié C, Saavedra H, Tróchez P, et al (1993) Life cycle, behaviour and natural enemies of *Liriomyza huidobrensis* (Blanchard) on bulb onion (*Allium cepa* L.). *Rev Colomb Entomol* 19:51–57
- International Plant Protection Convention (2017) Detection of *Liriomyza sativae* in Far North Queensland
- Iwasaki A (2004a) Effect of insecticides on *Liriomyza huidobrensis* (Blanchard)(Diptera: Agromyzidae) in Hokkaido [Japan]. *Annu Rep Soc Plant Prot North Japan*
- Iwasaki A (2004b) Effects of short photoperiod on cold hardiness and pupal period of the pea leafminer, *Liriomyza huidobrensis* (Blanchard) (Diptera: Agromyzidae)
- Kahinga, J. N., Gichuki, P. M., & Waigonjo, M. M. (2017). Managing passionfruit leafminer in the context of integrated-pest-management in the highland of central Kenya.
- Koch KC, Waterhouse DF (2000) Distribución e importancia de los artrópodos asociados a la agricultura y silvicultura en Chile. *ACIAR Monogr* 68:234 pp.
- Korytkowski C (1982) Contribución al Conocimiento de los Agromyzidae (Diptera: Acalyptratae) en el Perú. (Doctoral dissertation, Tesis (Magister Scientiae, en la especialidad de Entomología). Universidad Nacional Agraria La Molina, Programa Académico de Graduados)
- Korytkowski C (2014) Contribución al conocimiento de los Agromyzidae (Diptera: Muscomorpha) en el Perú. *Rev Peru Entomol* 49:1–106
- Kox LFF, Van Den Beld HE, Lindhout BI, De Goffau LJW (2005) Identification of economically important *Liriomyza* species by PCR-RFLP analysis. *EPPO Bull* 35:79–85. <https://doi.org/10.1111/j.1365-2338.2005.00807.x>
- Kuhnke KH, Wulfert I, Opitz B (1998) Zur Überwinterungsfähigkeit der Minierfliege *Liriomyza huidobrensis* im Freiland. *Gesunde Pflanz* 50:129–132
- Larriain PS, Munoz CM (1997) Seasonal abundance, alternative hosts, and parasitism on leafminer *Liriomyza huidobrensis* (Blanchard) (Diptera: Agromyzidae) affecting potato crops of the IV region of Chile. *Agric Tec* 57:290–296
- Larriain PS, Salas CF, Jana AC, Graña SF (2013) Major artichoke pests (*Cynara cardunculus* sub. *scolymus* L.) in coquimbo region, Chile. *Acta Horti* 983:301–306
- Larriain SP (2002) Insect and mite pest incidence on sweet pepinos *Solanum muricatum* (Ait.) cultivated in the IV Region, Chile. *Agricultura-Técnica* 62:15–26
- Macdonald I W, Reaser JK, Bright C, et al (2003) Invasive Alien Species in Southern Africa. *Glob Invasive*

- Maharjan R, Oh HW, Jung C (2014) Morphological and genetic characteristics of *Liriomyza huidobrensis* (Blanchard) (Diptera: Agromyzidae) infesting potato crops in Korea. J Asia Pac Entomol 17:281–286. <https://doi.org/10.1016/j.aspen.2014.01.013>
- Martin AD, Stanley-Horn D, Hallett RH (2005) Adult Host Preference and Larval Performance of *Liriomyza huidobrensis* (Diptera: Agromyzidae) on Selected Hosts. Environ Entomol 34:1170–1177. [https://doi.org/10.1603/0046-225x\(2005\)034\[1170:ahpalp\]2.0.co;2](https://doi.org/10.1603/0046-225x(2005)034[1170:ahpalp]2.0.co;2)
- Masetti A, Luchetti A, Mantovani B, Burgio G (2006) Polymerase Chain Reaction-Restriction Fragment Length Polymorphism Assays to Distinguish *Liriomyza huidobrensis* (Diptera: Agromyzidae) from Associated Species on Lettuce Cropping Systems in Italy. J Econ Entomol 99:1268–1272. <https://doi.org/10.1603/0022-0493-99.4.1268>
- Moura AP, Filho MM, Guimarães JA, Liz RS (2014) Manejo integrado de pragas do tomateiro para processamento industrial. Embrapa 1415–3033
- Mujica N (2007) Malezas hospederas de moscas minadoras (Diptera: Agromyzidae) y sus parasitoides en el ecosistema de papa en La Molina. Tesis Mg. Se. Lima, Perú. UNALM
- Mujica N, Kroschel J (2011) Leafminer Fly (Diptera: Agromyzidae) Occurrence, Distribution, and Parasitoid Associations in Field and Vegetable Crops Along the Peruvian Coast. Environ Entomol 40:217–230. <https://doi.org/10.1603/en10170>
- Musundire R, Chabi-Olaye A, Krüger K (2012) Host plant effects on morphometric characteristics of *Liriomyza huidobrensis*, *L. sativae* and *L. trifolii* (Diptera: Agromyzidae). J Appl Entomol 136:97–108. <https://doi.org/10.1111/j.1439-0418.2010.01597.x>
- Musundire R, Chabi-Olaye A, Löhr B, Krüger K (2011) Diversity of agromyzidae and associated hymenopteran parasitoid species in the afrotropical region: Implications for biological control. BioControl 56:1–9. <https://doi.org/10.1007/s10526-010-9312-z>
- Nakamura S, Masuda T, Mochizuki A, et al (2013) Primer design for identifying economically important *Liriomyza* species (Diptera: Agromyzidae) by multiplex PCR. Mol Ecol Resour 13:96–102. <https://doi.org/10.1111/1755-0998.12025>
- Niño L, Prieto L, Santiago V, Acevedo E (2009) Fluctuación poblacional de la mosca minadora (*Liriomyza huidobrensis* Blanchard) en cultivos de papa de Pueblo Llano, Estado Mérida, Venezuela. Entomotropica 24:65–70
- Noujeim E, Sakr J, Nemer N, others (2013) Potential of entomopathogenic nematodes application against *Liriomyza huidobrensis* Blanchard in Lebanon. In: Proceedings, 4th International Scientific Symposium Agrosys Bosnia and Herzegovina, Faculty of Agriculture, University of East Sarajevo. Bosnia and Herzegovina. pp 692–698
- Okoth CA, Deng AL, Tabu IM, et al (2014) Effect of Host Plant on Feeding, Biological and Morphological Parameters of *Liriomyza huidobrensis* Blanchard (Diptera: Agromyzidae). African Entomol 22:577–588. <https://doi.org/10.4001/003.022.0315>
- Olivera CR, Bordat D (1996) Influence of *Liriomyza* species (Diptera: Agromyzidae) and their host plants, on oviposition by *Opius dissitus* females (Hymenoptera: Braconidae). Ann Appl Biol 128:399–404. <https://doi.org/10.1111/j.1744-7348.1996.tb07101.x>
- Osorio A, Torres A, De La Riva F (1983) Producción de tomate (*Lycopersicon esculentum*, Mill.) regado por goteo, surcos y "caracol", en el Valle de Azapa1. IDESIA (Chile) 7:
- Pang B, Gao J, Zhou X, Wang J (2006) Relationship between host plant preference of *Liriomyza huidobrensis* (Blanchard)(Diptera: Agromyzidae) and secondary plant compounds and trichomes of host foliage. Kun chong xue bao Acta Entomol Sin 49:810–815
- Parish JB, Carvalho GA, Ramos RS, et al (2017) Host range and genetic strains of leafminer flies (Diptera: Agromyzidae) in eastern Brazil reveal a new divergent clade of *Liriomyza sativae*. 19:235–244. <https://doi.org/10.1111/afe.12202>
- Rauf A, Shepard BM, Johnson MW (2000) Leafminers in vegetables, ornamental plants and weeds in Indonesia: Surveys of host crops, species composition and parasitoids. Int J Pest Manag 46:257–266. <https://doi.org/10.1080/09670870050206028>

- Ripa SR, Rojas PS, Velasco G (1995) Releases of biological control agents of insect pests on Easter Island (Pacific Ocean). *Entomophaga* 40:427–440. <https://doi.org/10.1007/BF02373730>
- Romero Zuniga H, Zebisch T, Carballo M (1991) Life cycle and feeding preference of *Liriomyza huidobrensis* Blanchard (Diptera: Agromyzidae) on potato, celery and five important weeds in Cartago, Costa Rica. *Manejo Integr Plagas* 1–4
- Saethre M (1996) Pest risk assessment (PRA) for the South American leafminer *Liriomyza huidobrensis*
- Salvo A, Valladares G (2002) Plant-Related Intraspecific Size Variation in Parasitoids (Hymenoptera: Parasitica) of a Polyphagous Leafminer (Diptera: Agromyzidae). *Environ Entomol* 31:874–879. <https://doi.org/10.1603/0046-225x-31.5.874>
- Salvo A, Valladares GR (1997) An analysis of leaf-miner and plant host ranges of three *Chrysocharis* species (Hym.: Eulophidae) from Argentina. *Entomophaga* 42:417–426. <https://doi.org/10.1007/BF02769835>
- Saunders JL, Coto DT, King ABS (1998) Plagas invertebradas de cultivos anuales alimenticios en America Central
- Scheffer SJ, Lewis ML, Joshi RC (2006) DNA Barcoding Applied to Invasive Leafminers (Diptera: Agromyzidae) in the Philippines. *Ann Entomol Soc Am* 99:204–210. [https://doi.org/10.1603/0013-8746\(2006\)099\[0204:dbatil\]2.0.co;2](https://doi.org/10.1603/0013-8746(2006)099[0204:dbatil]2.0.co;2)
- Scheffer SJ, Wijesekara A, Visser D, Hallett RH (2001) Polymerase chain reaction-restriction fragment-length polymorphism method to distinguish *Liriomyza huidobrensis* from *L. langei* (Diptera: Agromyzidae) applied to three recent leafminer invasions. *J Econ Entomol* 94:1177–1182. <https://doi.org/10.1603/0022-0493-94.5.1177>
- Shepard BM, Samsudin, Braun AR (1998) Seasonal incidence of *Liriomyza huidobrensis* (Diptera: Agromyzidae) and its parasitoids on vegetables in Indonesia. *Int J Pest Manag* 44:43–47. <https://doi.org/10.1080/096708798228518>
- Shiao S, Wu W (2000) *Liriomyza huidobrensis* (Blanchard), a Newly Invaded Insect of Economic Importance to Taiwan (Diptera: Agromyzidae). 249–254
- Silva O (1993) Identification of the parasitoid complex on vegetable leaf miner *Liriomyza huidobrensis* (Blanchard) (Diptera: Agromyzidae) and some aspects of the biology of *Euparacrias phytomyzae* (Brethes) (Hymenoptera: Eulophidae).
- Sivapragasam A, Syed AR (1999) The problem and management of agromyzid leafminers on vegetables in Malaysia. 36–41
- Sivapragasam A, Syed AR, La Salle J, Ruwaida M (1999) Parasitoids of invasive agromyzid leafminers on vegetables in Peninsular Malaysia. In: *Proceedings of Symposium on Biological Control in the Tropics*. pp 8–19
- Spencer KA (1973) Agromyzidae (Diptera) of economic importance. 9:418. <https://doi.org/10.1007/978-94-017-0683-4>
- Spencer KA (1983) Leaf Mining Agromyzidae (Diptera) in Costa Rica. *Rev Biol Trop* 31:41–67. <https://doi.org/10.15517/rbt.v31i1.25127>
- Takano SI, Iwaizumi R, Nakanishi Y, Someya H (2008) Laboratory hybridization between the two clades of *Liriomyza huidobrensis* (Diptera: Agromyzidae). *Appl Entomol Zool* 43:397–402. <https://doi.org/10.1303/aez.2008.397>
- Tran DH (2009) Agromyzid leafminers and their parasitoids on vegetables in central Vietnam. *J ISSAAS* 15:21–33
- Ueno T (2006) Current status of insect pests attacking green bunching onion in central and southern Vietnam. *J Fac Agric Kyushu Univ* 51:275–283
- Valladares (1984) Sobre el Género *Liriomyza* Mik 1894 (Diptera, Agromyzidae) en la República Argentina. *rev soc ent argentina* 43:13–36
- Valladares G, Pinta D, Salvo A (1996) La mosca minadora *Liriomyza huidobrensis* Diptera, Agromyzidae en cultivos de Córdoba. *Hortic Argentina* 15:13–18
- Valladares G, Salvo A, Saini E (2011) Moscas minadoras del girasol y sus enemigos naturales. *RIA Rev Investig Agropecu* 37:180–188

- Valladares GR, Salvo A, Videla M (1999) Moscas minadoras en cultivos de Argentina. *Hortic Argentina* 18:44–45
- Verjel Manzano L, Mejia Florez M (2000) Desarrollo de los sistemas de producción cebolla de bulbo (Ocañera) y frijol rosado o Zaragoza mediante una agricultura orgánica libre de agroquímicos
- Wei J, Zou L, Kuang R, He L (2000) Influence of leaf tissue structure on host feeding selection by pea leafminer *Liriomyza huidobrensis* (Diptera: Agromyzidae). *Zool Stud* 39:295–300
- Weintraub P, Scheffer S, Visser D, et al (2017) The Invasive *Liriomyza huidobrensis* (Diptera: Agromyzidae): Understanding Its pest status and management globally. *J Insect Science* 17:. <https://doi.org/10.1093/jisesa/iew121>
- Weintraub PG (1999) Effects of cyromazine and abamectin on the leafminer, *Liriomyza huidobrensis* and its parasitoid, *Diglyphus isaea* in celery. *Ann Appl Biol* 135:547–554. <https://doi.org/10.1111/j.1744-7348.1999.tb00886.x>
- Wijesekara, A. (2010): Invasive alien species of agricultural importance in Sri Lanka: Have we managed them properly? In: *Invasive Alien Species in Sri Lanka – Strengthening Capacity to Control Their Introduction and Spread* (Eds: Marambe, B., Silva, P., Wijesundara, S. and Atapattu, N.), pp 115-122. Biodiversity Secretariat of the Ministry of Environment, Sri Lanka.
- Yábar E, Gianoli E, Echegaray ER (2002) Insect pests and natural enemies in two varieties of quinoa (*Chenopodium quinoa*) at Cusco, Peru. *J Appl Entomol* 126:275–280. <https://doi.org/10.1046/j.1439-0418.2002.00664.x>
- Yabaş C, Civelek HS, Ulubılır A (1995) The new leaf miner, *Liriomyza huidobrensis* (Blanchard 1926) on vegetables for Türkiye. *Türkiye Entomoloji Derg* 19:



Appendix 2: Leaf mining damage caused by established species







Table 16 provides a summary of established (native or naturalised) leaf mining species, their host overlap with pea leafminer and features that can be used to differentiate between serpentine leafminer and established leafminers.





It should be noted that because host plant records are poor for many native or naturalised species, there could be many more affected hosts than included here. The host overlap between serpentine leafminer and several native leafminer underscores the importance of sample collection and molecular diagnostics.

Figure 8 provides some general guidelines for differentiating between Dipteran and Lepidopteran leafminer larvae.

Table 16 Leaf mining damage caused by Australian dipterans or lepidopterans on leaves of cultivated crops and common weeds of cultivated crops

LEAFMINER	DISTINGUISHING FEATURES FROM SERPENTINE LEAFMINER DAMAGE	HOSTS (THOSE SHARED WITH SERPENTINE LEAFMINER ARE UNDERLINED)	PHOTO OF LEAF MINE
<i>Liriomyza brassicae</i>	None (all life stages are morphologically indistinguishable in appearance, without dissection, and behaviour from serpentine leafminer)	<u>Brassicaceae</u> <u>Capparaceae</u> <u>Tropaeolaceae</u> <u>Fabaceae</u>	
<i>Liriomyza chenopodii</i>	None (all life stages are morphologically indistinguishable in appearance, without dissection, and behaviour from serpentine leafminer)	<u>Amaranthaceae</u> <u>Caryophyllaceae</u> Any native hosts are as yet unrecorded (<i>L. chenopodii</i> is a native species)	

LEAFMINER	DISTINGUISHING FEATURES FROM SERPENTINE LEAFMINER DAMAGE	HOSTS (THOSE SHARED WITH SERPENTINE LEAFMINER ARE UNDERLINED)	PHOTO OF LEAF MINE
<i>Phytomyza syngenesiae</i>	<p>Pupation occurs inside the leaf mine.</p>  <p>Adults easily distinguishable from serpentine leafminer.</p>	<u>Asteracea</u> <u>Apiaceae</u> <u>Fabaceae</u>	
<i>Tropicomyia polyptya</i>	<p>Larvae are upper surface epidermal feeders (deeper parenchymatous tissues is not eaten). As such, mines have a silvery, film like appearance. Pupation occurs inside the mine.</p>  <p>Adults easily distinguishable from serpentine leafminer</p>	<p>Highly polyphagous Common hosts include plants in the <u>Passifloraceae</u>, <u>Euphorbiaceae</u>, <u>Fabaceae</u>, <u>Rubiaceae</u>, <u>Ruaceae</u>, and <u>Solanaceae</u> families, and more.</p>	
<i>Ophiomyia solanicola</i>	<p>Pupation occurs inside the leaf mine.</p>  <p>Adults easily distinguishable from serpentine leafminer</p>	<u>Solanaceae</u>	

LEAFMINER	DISTINGUISHING FEATURES FROM SERPENTINE LEAFMINER DAMAGE	HOSTS (THOSE SHARED WITH SERPENTINE LEAFMINER ARE UNDERLINED)	PHOTO OF LEAF MINE
<i>Ophiomyia alysicarpi</i>	<p>Mint green pupae can be found inside mines</p>  <p>Adults easily distinguishable from serpentine leafminer</p>	<i>Alysicarpus</i> sp. <i>Desmodium</i> sp.	
<i>Ophiomyia cornuta</i>	<p>Adults easily distinguishable from serpentine leafminer</p>	<u>Scaevola</u> sp. <u>Goodenia</u> sp.	
epidopteran leafminer	<p>Lepidopteran larvae can be distinguished from dipteran larvae on the spot, via a hand lens (Figure 10).</p> <p>For some lepidopteran species, when the larva is in its final stages, it forms a large blister at the end of the mine (see images on the right). Serpentine leafminer larvae do not create such blisters.</p>	Polyphagous (incl. <u>Fabaceae</u> , <u>Eucalypts</u> , native trees, etc)	

LEAFMINER	DISTINGUISHING FEATURES FROM SERPENTINE LEAFMINER DAMAGE	HOSTS (THOSE SHARED WITH SERPENTINE LEAFMINER ARE UNDERLINED)	PHOTO OF LEAF MINE
			




Figure 8 General guidelines for distinguishing dipteran vs lepidopteran leafminer larvae. Dipteran leafminer larvae (A) lack a head capsule and have a smooth body without clear segments. Lepidopteran leafminer larvae (B) have a distinct head capsule and are more visibly segmented. When attempting to distinguish a dipteran from lepidopteran leafminer larva on the spot, it is helpful to backlight the leaf as it is viewed under a hand lens, or to carefully excise the larva. Images: Elia Pirtle, Cesar Australia Pty Ltd

Surveillance for *L. huidobrensis* in Australia will be confounded by the presence of other leafmining insects already present in Australia, many of which create mines that are indistinguishable to a casual observer, and even in some cases indistinguishable to experts.

Table 17 lists genera that include hosts of *L. huidobrensis* but do not appear to contain of other Agromyzids within Australia or Australasia. It is important to note that host plant records for non-pest leafminer flies are highly incomplete, and some of the genera included herein could host leafminer flies that have not yet been reported. Host records for weeds and native plants are expected to be more incomplete than for cultivated crops and ornamentals, and despite poor host records for native leafminer. We can be more confident that native leafminer are not currently a notable presence within the genera containing cultivated crops and ornamentals, where sampling effort is expected to be higher. As such, leafmining damage detected within the genera included in these tables should be regarded as highly suspicious.

Error! Reference source not found. indicates where known overlap exists between hosts of *L. huidobrensis*, *L. trifolii* or *L. sativae* and native or naturalised agromyzids in Australasia.

Table 17 Host genera of *L. huidobrensis*, *L. sativae* or *L. trifolii* (including only records with comprehensive or partial evidence for lifecycle completion in the field, see Appendix 1) without any reported overlap with native or naturalised agromyzids already present in Australasia (see **Error! Reference source not found.** for known overlap). For each host, preferences of the three exotic *Liriomyza* are indicated (LH = *L. huidobrensis*; LT = *L. trifolii*; LS = *L. sativae*).

Genus	Family	Exotic <i>Liriomyza</i>	Host type
<i>Allium</i>	Amaryllidaceae	LH; LT; LS	Commercial
<i>Coriandrum</i>	Apiaceae	LH; LT; LS	Commercial
<i>Eryngium</i>	Apiaceae	LH	Commercial
<i>Petroselinum</i>	Apiaceae	LT	Commercial
<i>Colocasia</i>	Araceae	LH	Commercial
<i>Cichorium</i>	Asteraceae	LH	Commercial
<i>Crassocephalum</i>	Asteraceae	LH	Commercial
<i>Basella</i>	Basellaceae	LH	Commercial
<i>Eruca</i>	Brassicaceae	LH	Commercial
<i>Spinacea</i>	Chenopodiaceae	LH	Commercial
<i>Benincasa</i>	Cucurbitaceae	LH; LT; LS	Commercial
<i>Coccinia</i>	Cucurbitaceae	LT	Commercial
<i>Lagenaria</i>	Cucurbitaceae	LH; LT; LS	Commercial
<i>Luffa</i>	Cucurbitaceae	LH; LT; LS	Commercial
<i>Momordica</i>	Cucurbitaceae	LH; LT; LS	Commercial
<i>Sechium</i>	Cucurbitaceae	LH	Commercial
<i>Trichosanthes</i>	Cucurbitaceae	LT	Commercial
<i>Arachis</i>	Fabaceae	LT	Commercial
<i>Cicer</i>	Fabaceae	LH; LT; LS	Commercial
<i>Cyamopsis</i>	Fabaceae	LT	Commercial
<i>Lens</i>	Fabaceae	LS	Commercial
<i>Lotus</i>	Fabaceae	LH	Commercial
<i>Lupin</i>	Fabaceae	LH	Commercial
<i>Medicago</i>	Fabaceae	LH; LT; LS	Commercial
<i>Trigonella</i>	Fabaceae	LH; LT	Commercial
<i>Vicia</i>	Fabaceae	LH; LT; LS	Commercial
<i>Mentha</i>	Lamiaceae	LH; LS	Commercial
<i>Origanum</i>	Lamiaceae	LS	Commercial
<i>Linum</i>	Linaceae	LH	Commercial
<i>Abelmoschus</i>	Malvaceae	LH; LT; LS	Commercial
<i>Corchorus</i>	Malvaceae	LT	Commercial
<i>Gossypium</i>	Malvaceae	LT; LS	Commercial
<i>Hibiscus</i>	Malvaceae	LH; LT	Commercial
<i>Zizania</i>	Poaceae	LH	Commercial
<i>Lycium</i>	Solanaceae	LH	Commercial
<i>Thurnbergia</i>	Acanthaceae	LH	Ornamental
<i>Celosia</i>	Amaranthaceae	LH; LT	Ornamental
<i>Bupleurum</i>	Apiaceae	LH	Ornamental
<i>Centella</i>	Apiaceae	LH	Ornamental
<i>Oenanthe</i>	Apiaceae	LH	Ornamental
<i>Catharanthus</i>	Apocynaceae	LH	Ornamental
<i>Asclepias</i>	Asclepiadaceae	LT	Ornamental
<i>Chionodoxa</i>	Asparagaceae	LH	Ornamental
<i>Hemerocallis</i>	Asphodelaceae	LH	Ornamental
<i>Argyranthemum</i>	Asteraceae	LH	Ornamental

Genus	Family	Exotic <i>Liriomyza</i>	Host type
<i>Calendula</i>	Asteraceae	LH; LT; LS	Ornamental
<i>Centaurea</i>	Asteraceae	LH	Ornamental
<i>Conoclinium</i>	Asteraceae	LH	Ornamental
<i>Cosmos</i>	Asteraceae	LH	Ornamental
<i>Dimorphotheca</i>	Asteraceae	LT	Ornamental
<i>Echinacea</i>	Asteraceae	LT	Ornamental
<i>Eupatorium</i>	Asteraceae	LT	Ornamental
<i>Felicia</i>	Asteraceae	LS	Ornamental
<i>Flaveria</i>	Asteraceae	LT	Ornamental
<i>Kalimeris</i>	Asteraceae	LH	Ornamental
<i>Osteospermum</i>	Asteraceae	LH	Ornamental
<i>Pyrethrum</i>	Asteraceae	LH	Ornamental
<i>Tanacetum</i>	Asteraceae	LH	Ornamental
<i>Zinnia</i>	Asteraceae	LH; LT; LS	Ornamental
<i>Impatiens</i>	Balsaminaceae	LH	Ornamental
<i>Nasturtium</i>	Brassicaceae	LH; LS	Ornamental
<i>Calceolaria</i>	Calceolariaceae	LH	Ornamental
<i>Campanula</i>	Campanulaceae	LH	Ornamental
<i>Platycodon</i>	Campanulaceae	LH	Ornamental
<i>Trachelium</i>	Campanulaceae	LH; LT	Ornamental
<i>Dianthus</i>	Caryophyllaceae	LH; LT	Ornamental
<i>Gypsophila</i>	Caryophyllaceae	LH; LT; LS	Ornamental
<i>Bauhinia</i>	Fabaceae	LS	Ornamental
<i>Lathyrus</i>	Fabaceae	LH	Ornamental
<i>Lupinus</i>	Fabaceae	LH; LS	Ornamental
<i>Eustoma</i>	Gentianaceae	LH; LT	Ornamental
<i>Exacum</i>	Gentianaceae	LH	Ornamental
<i>Lisianthus</i>	Gentianaceae	LH; LT; LS	Ornamental
<i>Freesia</i>	Iridaceae	LH	Ornamental
<i>Gladiolus</i>	Iridaceae	LH	Ornamental
<i>Salvia</i>	Lamiaceae	LH; LT	Ornamental
<i>Lilium</i>	Liliaceae	LH	Ornamental
<i>Torenia</i>	Linderniaceae	LH	Ornamental
<i>Althaea</i>	Malvaceae	LH	Ornamental
<i>Moringa</i>	Moringaceae	LS	Ornamental
<i>Godetia</i>	Onagraceae	LH	Ornamental
<i>Oenothera</i>	Onagroidae	LH	Ornamental
<i>Dendrobium</i>	Orchidaceae	LS	Ornamental
<i>Papaver</i>	Papaveraceae	LH	Ornamental
<i>Antirrhinum</i>	Plantaginaceae	LH; LT; LS	Ornamental
<i>Myosotis</i>	Plumbaginaceae	LH	Ornamental
<i>Lagurus</i>	Poaceae	LH	Ornamental
<i>Phlox</i>	Polemoniaceae	LH; LT; LS	Ornamental
<i>Primula</i>	Primulaceae	LH	Ornamental
<i>Aquilegia</i>	Ranunculaceae	LS	Ornamental
<i>Delphinium</i>	Ranunculaceae	LH	Ornamental
<i>Nigella</i>	Ranunculaceae	LH	Ornamental
<i>Prunus</i>	Rosaceae	LS	Ornamental

Genus	Family	Exotic <i>Liriomyza</i>	Host type
<i>Rosa</i>	Rosaceae	LH; LT	Ornamental
<i>Diascia</i>	Scrophulariaceae	LH	Ornamental
<i>Viola</i>	Violaceae	LH	Ornamental
<i>Sagittaria</i>	Alismataceae	LH	Non cultivated
<i>Alternanthera</i>	Amaranthaceae	LT	Non cultivated
<i>Amaranthus</i>	Amaranthaceae	LH; LT; LS	Non cultivated
<i>Deeringia</i>	Amaranthaceae	LH	Non cultivated
<i>Gomphrena</i>	Amaranthaceae	LH	Non cultivated
<i>Iresine</i>	Amaranthaceae	LH	Non cultivated
<i>Alocasia</i>	Araceae	LH	Non cultivated
<i>Hydrocotyle</i>	Araliaceae	LH; LT; LS	Non cultivated
<i>Artemisia</i>	Asteraceae	LH; LT	Non cultivated
<i>Baccharis</i>	Asteraceae	LT	Non cultivated
<i>Carduus</i>	Asteraceae	LH	Non cultivated
<i>Dichrocephala</i>	Asteraceae	LH	Non cultivated
<i>Echinops</i>	Asteraceae	LH	Non cultivated
<i>Eclipta</i>	Asteraceae	LT	Non cultivated
<i>Emilia</i>	Asteraceae	LH	Non cultivated
<i>Erechtites</i>	Asteraceae	LH; LT	Non cultivated
<i>Galinsoga</i>	Asteraceae	LH; LT; LS	Non cultivated
<i>Helipterum</i>	Asteraceae	LH	Non cultivated
<i>Hemistepta</i>	Asteraceae	LH	Non cultivated
<i>Hymenopappus</i>	Asteraceae	LT	Non cultivated
<i>Lipochaeta</i>	Asteraceae	LS	Non cultivated
<i>Melanthera</i>	Asteraceae	LT	Non cultivated
<i>Parthenium</i>	Asteraceae	LT	Non cultivated
<i>Synedrella</i>	Asteraceae	LH; LT	Non cultivated
<i>Tridax</i>	Asteraceae	LT	Non cultivated
<i>Verbesina</i>	Asteraceae	LS	Non cultivated
<i>Xanthium</i>	Asteraceae	LT	Non cultivated
<i>Cordia</i>	Boraginaceae	LT	Non cultivated
<i>Capsella</i>	Brassicaceae	LH	Non cultivated
<i>Rorippa</i>	Brassicaceae	LH	Non cultivated
<i>Silene</i>	Caryophyllaceae	LH; LT	Non cultivated
<i>Malachium</i>	Caryophyllaceae	LH	Non cultivated
<i>Spergula</i>	Caryophyllaceae	LT	Non cultivated
<i>Commelina</i>	Commelinaceae	LT	Non cultivated
<i>Calystegia</i>	Convolvulaceae	LH	Non cultivated
<i>Convolvulus</i>	Convolvulaceae	LT	Non cultivated
<i>Pharbitis</i>	Convolvulaceae	LH	Non cultivated
<i>Ceratodes</i>	Cucurbitaceae	LS	Non cultivated
<i>Melothria</i>	Cucurbitaceae	LH	Non cultivated
<i>Melilotus</i>	Fabaceae	LH; LS	Non cultivated
<i>Poissonia</i>	Fabaceae	LS	Non cultivated
<i>Lamium</i>	Lamiaceae	LH	Non cultivated
<i>Leonurus</i>	Lamiaceae	LH	Non cultivated
<i>Melissa</i>	Lamiaceae	LS	Non cultivated
<i>Anoda</i>	Malvaceae	LS	Non cultivated

Genus	Family	Exotic <i>Liriomyza</i>	Host type
<i>Malva</i>	Malvaceae	LH; LS	Non cultivated
<i>Sida</i>	Malvaceae	LH; LS	Non cultivated
<i>Oxalis</i>	Oxalidaceae	LH; LT	Non cultivated
<i>Piriqueta</i>	Passifloraceae	LT	Non cultivated
<i>Linaria</i>	Plantaginaceae	LT	Non cultivated
<i>Veronica</i>	Plantaginaceae	LH	Non cultivated
<i>Limonium</i>	Plumbaginaceae	LH	Non cultivated
<i>Polygala</i>	Polygalaceae	LT	Non cultivated
<i>Rumex</i>	Polygonaceae	LH	Non cultivated
<i>Cardiospermum</i>	Sapindaceae	LT	Non cultivated
<i>Nemesia</i>	Scrophulariaceae	LH	Non cultivated
<i>Datura</i>	Solanaceae	LH; LT; LS	Non cultivated
<i>Nicotiana</i>	Solanaceae	LH; LT	Non cultivated
<i>Kallstroemia</i>	Zygophyllaceae	LT	Non cultivated
<i>Tribulus</i>	Zygophyllaceae	LT	Non cultivated

Table 18 Host genera of either *L. huidobrensis*, *L. sativae* or *L. trifolii* with known overlap with native or naturalised agromyzids already present in Australasia. Some of these species are stem miner, which should be straightforward to distinguish from the leaf mining damage caused by *L. huidobrensis*, *L. sativae* or *L. trifolii*. Records of *L. sativae* host plants within the Torres Strait Islands (TSI) and on the Cape York Peninsula (CYP) are included.

Host genus	Host family	Australasian Agromyzid genera	Sources
<i>Beta</i>	Amaranthaceae	<i>Liriomyza</i>	Benavent Corai et al. (2005); Spencer (1973); Spencer (1977); Spencer (1990)
<i>Chenopodium</i>	Amaranthaceae	<i>Liriomyza</i>	Benavent Corai et al. (2005); Spencer (1977); Spencer (1990)
<i>Spinacia</i>	Amaranthaceae	<i>Liriomyza</i>	Benavent Corai et al. (2005); Spencer (1973); Spencer (1977)
<i>Apium</i>	Apiaceae	<i>Melanagromyza</i>	Benavent Corai et al. (2005); Spencer (1973); Spencer (1977)
<i>Daucus</i>	Apiaceae	<i>Phytomyza</i>	Spencer (1973)
<i>Asparagus</i>	Asparagaceae	<i>Hexomyza</i>	Benavent Corai et al. (2005)
<i>Ageratum</i>	Asteraceae	<i>Melanagromyza</i>	Spencer (1977)
<i>Arctium</i>	Asteraceae	<i>Phytomyza</i>	Benavent Corai et al. (2005)
<i>Aster</i>	Asteraceae	<i>Calycomyza</i> , <i>Phytomyza</i>	Benavent Corai et al. (2005); Spencer (1977)
<i>Bellis</i>	Asteraceae	<i>Calycomyza</i> , <i>Phytomyza</i>	Benavent Corai et al. (2005)
<i>Bidens</i>	Asteraceae	<i>Melanagromyza</i>	Spencer (1977)
<i>Callistephus</i>	Asteraceae	<i>Calycomyza</i> , <i>Phytomyza</i>	Benavent Corai et al. (2005)
<i>Carthamus</i>	Asteraceae	<i>Melanagromyza</i>	Benavent Corai et al. (2005)
<i>Chrysanthemum</i>	Asteraceae	<i>Ophiomyia</i> , <i>Phytomyza</i> , <i>Melanagromyza</i>	Benavent Corai et al. (2005); Spencer (1977)
<i>Cineraria</i>	Asteraceae	<i>Phytomyza</i>	Benavent Corai et al. (2005)
<i>Cirsium</i>	Asteraceae	<i>Phytomyza</i>	Spencer (1977)
<i>Conyza</i>	Asteraceae	<i>Calycomyza</i> , <i>Phytomyza</i>	Benavent Corai et al. (2005)
<i>Coreopsis</i>	Asteraceae	<i>Phytomyza</i>	Benavent Corai et al. (2005); Spencer (1977)
<i>Craspedia</i>	Asteraceae	<i>Liriomyza</i>	Benavent Corai et al. (2005)
<i>Crepis</i>	Asteraceae	<i>Phytomyza</i>	Benavent Corai et al. (2005)
<i>Cynara</i>	Asteraceae	<i>Phytomyza</i>	Spencer (1973)
<i>Dahlia</i>	Asteraceae	<i>Phytomyza</i>	Benavent Corai et al. (2005)
<i>Erigeron</i>	Asteraceae	<i>Calycomyza</i>	Benavent Corai et al. (2005); Spencer (1977)
<i>Gaillardia</i>	Asteraceae	<i>Phytomyza</i>	Benavent Corai et al. (2005)
<i>Gerbera</i>	Asteraceae	<i>Phytomyza</i>	Benavent Corai et al. (2005)
<i>Gnaphalium</i>	Asteraceae	<i>Melanagromyza</i> , <i>Malanagromyza</i>	Benavent Corai et al. (2005); Spencer (1977)
<i>Helianthus</i>	Asteraceae	<i>Calycomyza</i> , <i>Phytomyza</i>	Benavent Corai et al. (2005)
<i>Helichrysum</i>	Asteraceae	<i>Liriomyza</i> , <i>Melanagromyza</i> , <i>Phytomyza</i>	Benavent Corai et al. (2005); Spencer (1977)
<i>Lactuca</i>	Asteraceae	<i>Liriomyza</i> *, <i>Phytomyza</i>	Benavent Corai et al. (2005); Present in TSI/CYP; Spencer (1973)
<i>Leucanthemum</i>	Asteraceae	<i>Phytomyza</i>	Benavent Corai et al. (2005)
<i>Picris</i>	Asteraceae	<i>Phytomyza</i>	Benavent Corai et al. (2005)
<i>Senecio</i>	Asteraceae	<i>Melanagromyza</i> , <i>Phytomyza</i>	Benavent Corai et al. (2005); Spencer (1977); Spencer (1977)
<i>Solidago</i>	Asteraceae	<i>Calycomyza</i>	Benavent Corai et al. (2005)
<i>Sonchus</i>	Asteraceae	<i>Phytomyza</i>	Benavent Corai et al. (2005); Spencer (1977)

Host genus	Host family	Australasian Agromyzid genera	Sources
<i>Tagetes</i>	Asteraceae	<i>Liriomyza</i> *	Present in TSI/CYP
<i>Taraxacum</i>	Asteraceae	<i>Phytomyza</i>	Benavent Corai et al. (2005)
<i>Tecoma</i>	Bignoniaceae	<i>Liriomyza</i> *	Present in TSI/CYP
<i>Barbarea</i>	Brassicaceae	<i>Liriomyza</i>	Benavent Corai et al. (2005)
<i>Brassica</i>	Brassicaceae	<i>Liriomyza</i> *	Benavent Corai et al. (2005); Present in TSI/CYP; Spencer (1977)
<i>Cardamine</i>	Brassicaceae	<i>Liriomyza</i>	Spencer (1977)
<i>Diplotaxis</i>	Brassicaceae	<i>Liriomyza</i>	Spencer (1977)
<i>Hirschfeldia</i>	Brassicaceae	<i>Liriomyza</i>	Benavent Corai et al. (2005)
<i>Matthiola</i>	Brassicaceae	<i>Liriomyza</i>	Benavent Corai et al. (2005); Spencer (1977)
<i>Raphanus</i>	Brassicaceae	<i>Liriomyza</i>	Benavent Corai et al. (2005); Present in TSI/CYP; Spencer (1977)
<i>Silene</i>	Caryophyllaceae	<i>Liriomyza</i>	Benavent Corai et al. (2005); Spencer (1977); Spencer (1990)
<i>Stellaria</i>	Caryophyllaceae	<i>Liriomyza</i>	Benavent Corai et al. (2005); Spencer (1977); Spencer (1990)
<i>Cleome</i>	Cleomaceae	<i>Liriomyza</i> *	Benavent Corai et al. (2005); Present in TSI/CYP
<i>Ipomoea</i>	Convolvulaceae	<i>Melanagromyza</i>	Benavent Corai et al. (2005)
<i>Citrullus</i>	Cucurbitaceae	<i>Liriomyza</i> *	Present in TSI/CYP
<i>Cucumis</i>	Cucurbitaceae	<i>Liriomyza</i> *	Present in TSI/CYP
<i>Cucurbita</i>	Cucurbitaceae	<i>Liriomyza</i> *	Present in TSI/CYP
<i>Euphorbia</i>	Euphorbiaceae	<i>Tropicomyia</i>	Spencer (1977)
<i>Ricinus</i>	Euphorbiaceae	<i>Liriomyza</i> *	Present in TSI/CYP
<i>Cajanus</i>	Fabaceae	<i>Melanagromyza</i> , <i>Ophiomyia</i>	Spencer (1973); Spencer (1977)
<i>Canavalia</i>	Fabaceae	<i>Ophiomyia</i>	Spencer (1973)
<i>Cassia</i>	Fabaceae	<i>Tropicomyia</i> , <i>Ophiomyia</i>	Spencer (1977); Spencer (1990)
<i>Crotalaria</i>	Fabaceae	<i>Liriomyza</i> *, <i>Ophiomyia</i>	Present in TSI/CYP; Spencer (1973); Spencer (1977)
<i>Dolichos</i>	Fabaceae	<i>Ophiomyia</i>	Spencer (1973); Spencer (1977)
<i>Glycine</i>	Fabaceae	<i>Melanagromyza</i> , <i>Ophiomyia</i>	Benavent Corai et al. (2005); Spencer (1973); Spencer (1977)
<i>Indigofera</i>	Fabaceae	<i>Liriomyza</i> *, <i>Ophiomyia</i>	Benavent Corai et al. (2005); Present in TSI/CYP; Spencer (1977); Spencer (1990)
<i>Lablab</i>	Fabaceae	<i>Ophiomyia</i>	Spencer (1973)
<i>Macroptilium</i>	Fabaceae	<i>Liriomyza</i> *, <i>Tropicomyia</i>	Present in TSI/CYP; Spencer (1977)
<i>Millettia</i>	Fabaceae	Unknown	Pirtle pers. comm.
unknown	Fabaceae	<i>Melanagromyza</i>	Spencer (1977)
<i>Phaseolus</i>	Fabaceae	<i>Ophiomyia</i> , <i>Tropicomyia</i> , <i>Melanagromyza</i> , <i>Ophiomyia</i>	Spencer (1973); Spencer (1977)
<i>Pisum</i>	Fabaceae	<i>Liriomyza</i> , <i>Phytomyza</i> , <i>Tropicomyia</i> , <i>Ophiomyia</i>	Benavent Corai et al. (2005); Spencer (1973); Spencer (1977)
<i>Senna</i>	Fabaceae	<i>Tropicomyia</i>	Spencer (1977)
<i>Sesbania</i>	Fabaceae	<i>Liriomyza</i> *	Present in TSI/CYP
<i>Trifolium</i>	Fabaceae	<i>Ophiomyia</i>	Benavent Corai et al. (2005); Spencer (1977)
<i>Vigna</i>	Fabaceae	<i>Liriomyza</i> *, <i>Melanagromyza</i> , <i>Tropicomyia</i> , <i>Ophiomyia</i>	Present in TSI/CYP; Pirtle et al. (2020); Spencer (1973); Spencer (1977); Spencer 1977

Host genus	Host family	Australasian Agromyzid genera	Sources
<i>Scaevola</i>	Goodeniaceae	<i>Liriomyza</i> , <i>Ophiomyia</i>	Benavent Corai et al. (2005); Pirtle pers. comm.; Spencer (1977)
<i>Hydrangea</i>	Hydrangeaceae	<i>Tropicomyia</i>	Spencer (1977)
<i>Ocimum</i>	Lamiaceae	<i>Liriomyza</i> *	Pirtle et al. (2020)
<i>Stephania</i>	Menispermaceae	<i>Tropicomyia</i>	Spencer (1977); Spencer (1990)
<i>Passiflora</i>	Passifloraceae	<i>Tropicomyia</i> , <i>Liriomyza</i> *	Pirtle et al. (2020); Spencer (1977); Spencer (1990)
<i>Turnera</i>	Passifloraceae	<i>Liriomyza</i> *	Pirtle et al. (2020)
<i>Plantago</i>	Plantaginaceae	<i>Liriomyza</i> , <i>Phytomyza</i>	Benavent Corai et al. (2005); Spencer (1977);
<i>Hordeum</i>	Poaceae	<i>Ceradontha</i> , <i>Pseudonapomyza</i>	Benavent Corai et al. (2005); Spencer (1973)
<i>unknown</i>	Poaceae	<i>Agromyza</i> , <i>Ceradontha</i>	Spencer (1973)
<i>Setaria</i>	Poaceae	<i>Agromyza</i>	Benavent Corai et al. (2005)
<i>Triticum</i>	Poaceae	<i>Ceradontha</i> , <i>Pseudonapomyza</i>	Benavent Corai et al. (2005); Spencer (1973)
<i>Zea</i>	Poaceae	<i>Pseudonapomyza</i>	Spencer (1973)
<i>Portulaca</i>	Portulacaceae	<i>Liriomyza</i>	Benavent Corai et al. (2005)
<i>Ranunculus</i>	Ranunculaceae	<i>Phytomyza</i> , <i>Napomyza</i>	Spencer (1977); Spencer (1990)
<i>Passiflora</i>	Resedaceae	<i>Tropicomyia</i>	Spencer (1990)
<i>Coffea</i>	Rubiaceae	<i>Tropicomyia</i>	Spencer (1973); Spencer (1977)
<i>Citrus</i>	Rutaceae	<i>Tropicomyia</i>	Spencer (1973); Spencer (1977)
<i>Capsicum</i>	Solanaceae	<i>Tropicomyia</i> , <i>Liriomyza</i> *	Pirtle et al. (2020); Spencer (1977)
<i>Cestrum</i>	Solanaceae	<i>Tropicomyia</i>	Spencer (1977)
<i>Lycopersicum</i>	Solanaceae	<i>Liriomyza</i> *	Present in TSI/CYP
<i>Petunia</i>	Solanaceae	<i>Liriomyza</i> *	Present in TSI/CYP
<i>Physalis</i>	Solanaceae	<i>Liriomyza</i> *	Present in TSI/CYP; Pirtle et al. (2020)
<i>Solanum</i>	Solanaceae	<i>Ophiomyia</i> , <i>Liriomyza</i> *	Pirtle pers. comm.; Present in TSI/CYP; Spencer (1977)
<i>Tropaeolum</i>	Tropaeolaceae	<i>Liriomyza</i>	Benavent Corai et al. (2005); Spencer (1977); Spencer (1990)
<i>Lantana</i>	Verbenaceae	<i>Calycomyza</i> , <i>Tropicomyia</i> , <i>Ophiomyia</i>	Benavent Corai et al. (2005); Spencer (1977)
<i>Stachytarpheta</i>	Verbenaceae	<i>Liriomyza</i> *	Pirtle et al. (2020)
<i>Verbena</i>	Verbenaceae	<i>Calycomyza</i>	Benavent Corai et al. (2005)

* Includes *Liriomyza sativae* within TSI/CYP

References

- Benavent-Corai, J., Martinez, M., & Peydró, R. J. (2005). Catalogue of the hosts-plants of the world Agromyzidae (Diptera). *Bolletino di Zoologia Agraria e di Bachicoltura. Serie II.* 37, 1–97.
- Pirtle, E., Van Rooyen, A., Maino, J., Weeks, A., & Umina, P. (2020). A molecular method for biomonitoring of an exotic plant-pest: Leafmining for environmental DNA. *Authorea Preprints*.
- Spencer, K. A. (1973). *Agromyzidae (Diptera) of economic importance (Vol. 9)*. Springer Science & Business Media.
- Spencer, K. A. (1977). A revision of the Australian Agromyzidae (Diptera). *Western Australian Museum, Special Publication*, 8, 1-255.
- Spencer, K. A. (1990). *Host specialization in the world Agromyzidae (Diptera) (Vol. 45)*. Springer Science & Business Media.

Appendix 3: Permits for the control of serpentine leafminer and other *Liriomyza* spp. on different host crops

Table 19 lists the agriculturally important hosts of serpentine leafminer (refer to Appendix 1) and notes where suitable pesticides are available and approved by the APVMA for control on each host crop. Blank cells denote where suitable pesticides have not yet been approved or identified. Refer to the APVMA permit database (<https://portal.apvma.gov.au/permits>) or the most up to date information on current leafminer permits.

Table 19 Insecticides control options for different crop groups

Relevant Crop group	Crop/host plant	Pesticide	Permit number(s)/ status	Target pest (as per permit)	Expiration date	Withholding information/notes	Suitable for management or eradication ²
Brassica vegetables	<u>Bok choy; pak-choy; turnip</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	Treated crop must be destroyed. Treated crops must not be made available for human consumption Do not graze or cut for stock food	Eradication only
Brassica vegetables	<u>Broccoli</u>	Emamectin benzoate	PER87563	<i>Liriomyza</i> spp. (including: <i>L. sativae</i>)	30 June 2024	3 days (harvest) Do not graze or cut for stock food	Management or eradication
Brassica vegetables	<u>Broccoli</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication

² Some pesticides will be more suitable for eradication than for the ongoing management of the pest. This is because either they will need to be used at high rates to be effective, there are potential resistance issues, or the pesticide is not likely to fit into existing IPM systems as well as alternative pesticides. Other chemicals provide good control of the pest and are also compatible with IPM systems and will be more suitable for the ongoing management of the pest. To highlight this, pesticides are noted for either eradication, or for management and eradication.

Brassica vegetables	<u>Brussels sprouts</u>	Emamectin benzoate	PER87563	<i>Liriomyza</i> spp. (including: <i>L. sativae</i>)	30 June 2024	3 days (harvest) Do not graze or cut for stock food	Management or eradication
Brassica vegetables	<u>Brussels sprouts</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	Treated crop must be destroyed. Treated crops must not be made available for human consumption Do not graze or cut for stock food	Eradication only
Brassica vegetables	<u>Cabbage</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Brassica vegetables	<u>Cabbage</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	Treated crop must be destroyed. Treated crops must not be made available for human consumption Do not graze or cut for stock food	Eradication only
Brassica vegetables	<u>Cabbage</u>	Emamectin benzoate	PER87563	<i>Liriomyza</i> spp. (including: <i>L. sativae</i>)	30 June 2024	3 days (harvest) Do not graze or cut for stock food	Management or eradication
Brassica vegetables	<u>Cauliflower</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	Treated crop must be destroyed. Treated crops must not be made available for human consumption Do not graze or cut for stock food	Eradication only
Brassica vegetables	<u>Cauliflower</u>	Emamectin benzoate	PER87563	<i>Liriomyza</i> spp. (including: <i>L. sativae</i>)	30 June 2024	3 days (harvest) Do not graze or cut for	Management or eradication

				<i>sativae</i>)		stock food	
Brassica vegetables	<u>Indian mustard</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	Treated crop must be destroyed. Treated crops must not be made available for human consumption Do not graze or cut for stock food	Eradication only
Bulb vegetables	<u>Garlic</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	14 days (harvest) Do not graze or cut for stock food	Management or eradication
Bulb vegetables	<u>Garlic</u>	Cyantraniliprole	PER90387	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 December 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Bulb vegetables	<u>Leek</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	14 days (harvest) Do not graze or cut for stock food	Management or eradication
Bulb vegetables	<u>Leek</u>	Cyantraniliprole	PER90387	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 December 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Bulb vegetables	<u>Onions</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	30 days (harvest) Do not graze or cut for stock food	Management or eradication
Bulb vegetables	<u>Onions</u>	Cyantraniliprole	PER90387	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 December 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication

Bulb vegetables	<u>Shallots</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	14 days (harvest) Do not graze or cut for stock food	Management or eradication
Bulb vegetables	<u>Shallots</u>	Cyantraniliprole	PER90387	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 December 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Bulb vegetables	<u>Spring onion</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	14 days (harvest) Do not graze or cut for stock food	Management or eradication
Bulb vegetables	<u>Spring onion</u>	Cyantraniliprole	PER90387	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 December 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Cereal grains	Barley	No suitable pesticides identified					
Fruit trees (non-bearing)	Any trees that will not bear fruit for the next 12 months	Cyromazine	PER83506	Larvae of leafminers	31 October 2022	Do not ship within 7 days of treatment	Management or eradication
Fruiting vegetables (Cucurbits – including melons)	<u>Angled luffa, sing-kwa</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (Cucurbits – including melons)	<u>Angled luffa, sing-kwa</u>	Cyantraniliprole	PER90387	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 December 2023	1 day (harvest) Do not graze or cut for stock food	Management or eradication

Fruiting vegetables (Cucurbits – including melons)	<u>Angled luffa, sing-kwa</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (Cucurbits – including melons)	<u>Bottle gourd</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (Cucurbits – including melons)	<u>Bottle gourd</u>	Cyantraniliprole	PER90387	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 December 2023	1 day (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (Cucurbits – including melons)	<u>Bottle gourd</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (Cucurbits – including melons)	<u>Cucumber</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (Cucurbits – including melons)	<u>Cucumber</u>	Cyantraniliprole	PER90387	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 December 2023	1 day (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting	<u>Cucumber</u>	Cyromazine	PER81867	<i>Liriomyza</i>	30 November	7 days (harvest)	Management or

vegetables (Cucurbits – including melons)				<i>sativae</i> and <i>L. huidobrensis</i>	2023	Do not graze or cut for stock food	eradication
Fruiting vegetables (Cucurbits – including melons)	<u>Melon (including honeydew, rock melon)</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (Cucurbits – including melons)	<u>Melon (including honeydew, rock melon)</u>	Cyantraniliprole	PER90387	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 December 2023	1 day (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (Cucurbits – including melons)	<u>Melon (including honeydew, rock melon)</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (Cucurbits – including melons)	<u>Pumpkin (e.g. butternut)</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (Cucurbits – including melons)	<u>Pumpkin (e.g. butternut)</u>	Cyantraniliprole	PER90387	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 December 2023	1 day (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (Cucurbits – including melons)	<u>Bottle gourd</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication

including melons)							
Fruiting vegetables (Cucurbits – including melons)	<u>Pumpkin (e.g. Queensland blue)</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (Cucurbits – including melons)	<u>Pumpkin (e.g. Queensland blue)</u>	Cyantraniliprole	PER90387	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 December 2023	1 day (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (Cucurbits – including melons)	<u>Pumpkin (e.g. Queensland blue)</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (Cucurbits – including melons)	<u>Smooth luffa, sponge gourd</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (Cucurbits – including melons)	<u>Smooth luffa, sponge gourd</u>	Cyantraniliprole	PER90387	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 December 2023	1 day (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (Cucurbits – including melons)	<u>Smooth luffa, sponge gourd</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting	<u>Watermelon</u>	Abamectin	PER81876	<i>Liriomyza</i>	30 April 2024	7 days (harvest)	Management or

vegetables (Cucurbits – including melons)				<i>sativae</i> and <i>L. huidobrensis</i>		Do not graze or cut for stock food	eradication
Fruiting vegetables (Cucurbits – including melons)	<u>Watermelon</u>	Cyantranilprole	PER90387	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 December 2023	1 day (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (Cucurbits – including melons)	<u>Watermelon</u>	Cyromazine	PER81867	<i>Liriomyza</i> <i>sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (Cucurbits – including melons)	<u>Wax gourd</u>	Abamectin	PER81876	<i>Liriomyza</i> <i>sativae</i> and <i>L. huidobrensis</i>	30 April 2024	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (Cucurbits – including melons)	<u>Wax gourd</u>	Cyantranilprole	PER90387	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 December 2023	1 day (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (Cucurbits – including melons)	<u>Wax gourd</u>	Cyromazine	PER81867	<i>Liriomyza</i> <i>sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication

Fruiting vegetables (Cucurbits – including melons)	<u>Zucchini</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (Cucurbits – including melons)	<u>Zucchini</u>	Cyrantraniliprole	PER90387	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 December 2023	1 day (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (Cucurbits – including melons)	<u>Zucchini</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (other than cucurbits)	Bell pepper	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (other than cucurbits)	Bell pepper	Cyrantraniliprole	PER90387	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 December 2023	1 day (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (other than cucurbits)	Bell pepper	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (other than cucurbits)	<u>Capsicum, chili*</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables	<u>Capsicum, chili*</u>	Cyrantraniliprole	PER90387	<i>Liriomyza</i> spp. (including:	31 December 2023	1 day (harvest) Do not graze or cut for	Management or eradication

(other than cucurbits)				<i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)		stock food	
Fruiting vegetables (other than cucurbits)	<u>Capsicum, chili*</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (other than cucurbits)	<u>Capsicum, chili*</u>	Spirotetramat	PER88640	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 May 2023	1 day (harvest) Do not graze or cut for stock food	Management
Fruiting vegetables (other than cucurbits)	<u>Eggplant</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (other than cucurbits)	<u>Eggplant</u>	Cyantranilprole	PER90387	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 December 2023	1 day (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (other than cucurbits)	<u>Eggplant</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (other than cucurbits)	<u>Eggplant</u>	Spirotetramat	PER88640	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 May 2023	1 day (harvest) Do not graze or cut for stock food	Management
Fruiting vegetables (other than cucurbits)	<u>Okra*</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	7 days (harvest) Do not graze or cut for stock food	Management or eradication

Fruiting vegetables (other than cucurbits)	<u>Okra*</u>	Cyantraniliprole	PER90387	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 December 2023	1 day (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (other than cucurbits)	<u>Okra*</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (other than cucurbits)	<u>Tomato</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (other than cucurbits)	<u>Tomato</u>	Cyantraniliprole	PER90387	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 December 2023	1 day (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (other than cucurbits)	<u>Tomato</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Fruiting vegetables (other than cucurbits)	<u>Tomato</u>	Spirotetramat	PER88640	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 May 2023	1 day (harvest) Do not graze or cut for stock food	Management
Herbs	<u>Basil</u>	No suitable pesticides identified					
Herbs	<u>Oregano</u>	No suitable pesticides identified					

Herbs	<u>Parsley</u> ³	Spirotetramat	PER88640	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 May 2023	3 days (harvest) Do not graze or cut for stock food	Management
Head lettuce	<u>Lettuce (Head type only)</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Leafy vegetables (including brassica leafy vegetables)	<u>Lettuce</u>	Spirotetramat	PER88640	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 May 2023	1 day (harvest) Do not graze or cut for stock food	Management
Leafy vegetables (including brassica leafy vegetables)	<u>Spinach</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	14 days (harvest) Do not graze or cut for stock food	Management or eradication
Leafy vegetables (including brassica leafy vegetables)	<u>Spinach</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	Treated crop must be destroyed. Treated crops must not be made available for human consumption Do not graze or cut for stock food	Eradication only
Leafy vegetables	<u>Spinach and silverbeet</u>	Chlorantraniliprole	PER87631	<i>Liriomyza</i> spp. (including:	30 June 2024	3 days (harvest) 7 days (grazing or	Management or eradication

³ Note: Parsley is not a recorded host for *L. sativae* but may be a host of other *Liriomyza* spp. and is listed as one of the crops that the permit covers.

(including brassica leafy vegetables)				<i>L. sativae</i> , <i>L. brassicae</i> , <i>L. huidobrensis</i>)		cutting for stock food)	
Legume vegetables	<u>Common bean;</u> <u>French beans,</u> <u>kidney beans</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Legume vegetables	<u>Common bean;</u> <u>French beans,</u> <u>kidney beans</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Legume vegetables	<u>Common bean;</u> <u>French beans,</u> <u>kidney beans</u>	Spirotetramat	PER88640	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 May 2023	7 days (harvest) 7 days (grazing and hay)	Management
Legume vegetables	<u>Faba bean, broad bean</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Legume vegetables	<u>Faba bean, broad bean</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Legume vegetables	<u>Mung bean</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Legume vegetables	<u>Mung bean</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Legume vegetables	<u>Pea, snow peas,</u> <u>sugar snap peas,</u> <u>field pea</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Legume vegetables	<u>Pea, snow peas,</u> <u>sugar snap peas,</u> <u>field pea</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Legume vegetables	<u>Pea, snow peas,</u> <u>sugar snap peas,</u>	Spinetoram	PER87878	<i>Liriomyza</i> spp.	28 February 2023	3 days (harvest); 14 days (grazing and	Management or eradication

	<u>field pea</u>					hay) Do not allow dairy cattle to graze treated forage	
Legume vegetables	<u>Pea, snow peas, sugar snap peas, field pea</u>	Spirotetramat	PER88640	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 May 2023	3 days (harvest); 3 days (grazing and hay)	Management
Legume vegetables	Red bean	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Legume vegetables	Red bean	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Legume vegetables	<u>Snake bean</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Legume vegetables	<u>Snake bean</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Not classified	Field mustard	No suitable pesticides identified					
Not classified	<u>Lucerne</u>	No suitable pesticides identified					
Nursery stock (non-food)	Various non-food nursery plants (e.g. chrysanthemum etc.)	Cyromazine	PER83506	Larvae of leafminers	31 October 2022	Do not ship within 7 days of treatment	Management or eradication
Nursery stock	Seedlings, tubes	Abamectin	PER88977	Leafminers	30	Not required when	Management or

(non-food)	and plugs, potted colour, trees and shrubs, foliage plants, palms, grasses, fruiting plants (non-bearing), cut flowers and ornamentals.	Azadirachtin Cyromazine Emamectin Chlorantraniliprole + thiamethoxam Cyantraniliprole Indoxacarb Spinetoram		(<i>Liriomyza</i> spp.) including Vegetable leafminer (<i>Liriomyza sativae</i>)	November, 2022	used as directed	eradication
Oilseeds	<u>Cotton</u>	Spinetoram has been identified as suitable for the control of Vegetable leafminer and other <i>Liriomyza</i> spp. on cotton as overseas and Australian use patterns are comparable. However, no permit has been applied for this use to date					
Pulses	<u>Pulses</u>	Dimethoate	PER89184	<i>Liriomyza sativae</i> , <i>Liriomyza trifolii</i> , <i>Liriomyza huidobrensis</i>	31 March 2025	Harvest: Do not harvest for 14 days after application. Grazing: Do not graze or cut for stock food for 14 days after application.	Management
Root and tuber vegetables	<u>Beets (beetroot, silver beet, sugar beets)</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	14 days (harvest) Do not graze or cut for stock food	Management or eradication
Stalk and stem vegetables	<u>Beets (beetroot, silver beet, sugar beets)</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Root and tuber vegetables	<u>Carrot</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	14 days (harvest) Do not graze or cut for stock food	Management or eradication
Stalk and stem vegetables	<u>Carrot</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Root and tuber	<u>Potato</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and	30 April 2024	14 days (harvest) Do not graze or cut for	Management or eradication

vegetables				<i>L. huidobrensis</i>		stock food	
Root and tuber vegetables	<u>Potato</u>	Cyrantraniliprole	PER90387	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 December 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Stalk and stem vegetables	<u>Potato</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Root and tuber vegetables	<u>Radish</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	14 days (harvest) Do not graze or cut for stock food	Management or eradication
Stalk and stem vegetables	<u>Radish</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Stalk and stem vegetables	<u>Celery</u>	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	7 days (harvest); do not graze	Management or eradication
Stalk and stem vegetables	<u>Celery</u>	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Stalk and stem vegetables	<u>Celery</u>	Spirotetramat	PER88640	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 May 2023	3 days (harvest) Do not graze or cut for stock food	Management or eradication
Stalk and stem vegetables	Rhubarb ⁴	Abamectin	PER81876	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 April 2024	7 days (harvest); do not graze	Management or eradication

⁴ Note: Rhubarb is not a recorded host for *L. sativae* or *L. huidobrensis* but may be a host of other *Liriomyza* spp. and is part of the 'Stalk and Stem Vegetable Crop Group' that the permit covers.

Stalk and stem vegetables	Rhubarb ³	Cyromazine	PER81867	<i>Liriomyza sativae</i> and <i>L. huidobrensis</i>	30 November 2023	7 days (harvest) Do not graze or cut for stock food	Management or eradication
Stalk and stem vegetables	Rhubarb ³	Spirotetramat	PER88640	<i>Liriomyza</i> spp. (including: <i>L. sativae</i> , <i>L. trifolii</i> , <i>L. huidobrensis</i>)	31 May 2023	3 days (harvest) Do not graze or cut for stock food	Management or eradication

Appendix 4: Additional impact prediction outputs

Table 20 State level accumulated unmitigated impacts in millions of dollars after 3 years resulting from a spring (September) incursion of *L. huidobrensis* at key entry points across Australia. Simulations were replicated 10 times with means and standard deviations shown in parentheses. The proportion crop impact was fixed at 10% in order to explore variability due to incursion location and the size and distribution of different industries. Thus, host preferences of the pests do not influence the predicted crop impacts, and impacts of low preference hosts may be overestimated while impacts of high preference hosts may be underestimated. All cells with a value of 0.00 represent unmitigated impacts less than \$10,000.

Crop	INCURSION LOCATION															
	NSW			NT		QLD			SA	TAS	VIC		WA			
	Griffith	Sydney	Western Sydney	Darwin	Katherine	Bowen	Bundaberg	Toowoomba	Adelaide	Devonport	Ballarat	Melbourne	Albany	Carnarvon	Geraldton	Perth
Beans	0.00 (0.00)	0.00 (0.00)	0.05 (0.00)	0.00 (0.00)	0.00 (0.00)	1.32 (0.00)	1.50 (0.00)	0.03 (0.02)	0.00 (0.00)	1.17 (0.00)	0.02 (0.01)	0.05 (0.02)	0.01 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Broccoli	0.03 (0.00)	0.00 (0.00)	0.03 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.46 (0.01)	0.00 (0.00)	0.38 (0.00)	1.07 (1.08)	3.49 (1.63)	0.04 (0.00)	0.00 (0.00)	0.00 (0.00)	0.01 (0.01)
Brussels	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	2.41 (0.00)	0.32 (0.00)	0.02 (0.06)	0.03 (0.08)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Cabbages	0.00 (0.00)	0.02 (0.03)	1.20 (0.02)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.35 (0.01)	0.10 (0.00)	0.12 (0.00)	0.12 (0.12)	0.34 (0.16)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Capsicum	0.04 (0.00)	0.00 (0.00)	0.07 (0.00)	0.00 (0.00)	0.00 (0.00)	3.68 (0.00)	1.69 (0.00)	0.00 (0.01)	0.00 (0.00)	1.47 (0.00)	0.08 (0.21)	0.01 (0.02)	0.04 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Carrots	0.02 (0.00)	0.00 (0.01)	0.02 (0.01)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.07 (0.02)	0.00 (0.00)	3.72 (0.00)	0.05 (0.04)	0.03 (0.03)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.02 (0.03)
Cauliflowers	0.00 (0.00)	0.01 (0.01)	0.05 (0.01)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.44 (0.01)	0.00 (0.00)	0.97 (0.00)	0.57 (0.64)	2.00 (0.96)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.01)
Flowers	0.00 (0.00)	0.44 (0.48)	3.87 (0.48)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.88 (0.07)	0.53 (0.18)	0.64 (0.04)	0.63 (0.02)	1.42 (0.61)	0.47 (0.51)	0.12 (0.00)	0.00 (0.00)	0.00 (0.00)	0.01 (0.00)
Lettuces	0.00	0.05	0.61	0.00	0.00	0.00	0.07	1.47	0.00	0.01	0.97	2.00	0.00	0.00	0.00	0.02

INCURSION LOCATION																
	(0.00)	(0.05)	(0.05)	(0.00)	(0.00)	(0.00)	(0.00)	(0.02)	(0.00)	(0.00)	(0.56)	(0.91)	(0.00)	(0.00)	(0.00)	(0.02)
Melons	0.90 (0.00)	0.00 (0.00)	0.03 (0.00)	0.00 (0.00)	0.17 (0.00)	0.17 (0.00)	3.98 (0.00)	0.11 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Nurseries	0.89 (0.00)	0.63 (0.46)	5.47 (0.54)	0.00 (0.00)	0.00 (0.00)	0.29 (0.00)	0.97 (0.19)	3.94 (1.86)	2.55 (0.02)	0.66 (0.01)	4.16 (2.53)	1.95 (1.31)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.03 (0.01)
Onions	0.09 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.21 (0.01)	0.00 (0.00)	3.71 (0.00)	0.01 (0.01)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Peas	0.00 (0.00)	0.00 (0.00)	0.01 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.29 (0.00)	0.00 (0.00)	0.00 (0.00)	0.63 (0.00)	0.00 (0.00)	0.00 (0.00)	0.01 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Potatoes	2.14 (0.00)	0.01 (0.02)	0.32 (0.01)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	2.40 (0.00)	0.51 (0.01)	0.05 (0.00)	5.99 (0.00)	2.54 (0.29)	0.30 (0.37)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.01)
Pumpkins	0.63 (0.00)	0.00 (0.00)	0.04 (0.00)	0.00 (0.00)	0.00 (0.00)	0.31 (0.00)	0.35 (0.00)	0.20 (0.01)	0.00 (0.00)	0.13 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Tomatoes	0.25 (0.00)	0.02 (0.02)	0.92 (0.08)	0.00 (0.00)	0.00 (0.00)	9.11 (0.00)	5.50 (0.00)	0.29 (0.02)	0.00 (0.00)	0.34 (0.00)	0.18 (0.55)	0.00 (0.01)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Total	5.00 (0.00)	1.19 (1.01)	12.68 (1.08)	0.00 (0.00)	0.17 (0.00)	14.88 (0.00)	17.65 (0.24)	8.62 (2.09)	5.76 (0.06)	20.26 (0.03)	11.21 (3.35)	10.67 (4.44)	0.22 (0.00)	0.00 (0.00)	0.00 (0.00)	0.09 (0.08)

Figure 9 Australia wide accumulating unmitigated impacts in millions of dollars after 3 years resulting from a spring (September) incursion of *L. huidobrensis* at key entry and establishment points across Australia.

